



# IJEAST

INTERNATIONAL JOURNAL  
OF ENGINEERING APPLIED SCIENCE  
AND TECHNOLOGY



**VOLUME : 9    ISSUE : 03    Print / Issue Publication Date: 09-Oct-2024**



**ISSN : 2455-2143**



**DOI : 10.33564/IJEAST.2024.v09i03.014**

Indexed In



[WWW.IJEAST.COM](http://WWW.IJEAST.COM)

[editor@ijeast.com](mailto:editor@ijeast.com)

# STUDY ON KNOWLEDGE AND ITS ASSOCIATED FACTORS REGARDING ANEMIA AMONG WOMEN AGED 18-45 IN AN URBAN SLUM IN BANGALORE, KARNATAKA.

Geethanjali Jerald  
MSc (N), PGDHP  
Professor

Department of Community Health Nursing  
Christ College of Nursing, Mangalore, Karnataka.

Dr. Shantharani N  
MDS

Department of Pedodontics and Preventive Dentistry  
Bapuji Dental College and Hospital  
Davangere, Karnataka.

**Abstract:** Background of the study: Anemia is a significant health issue worldwide. Iron Deficiency Anemia is one of the most common forms of anemia. A cross-sectional study was carried out to evaluate the awareness and related factors among women aged 18-45 regarding anemia in an urban slum in Bangalore, Karnataka.

**Methods:** A cross-sectional descriptive research approach was adopted in this study. The systematic random sampling technique was used to select 200 women of reproductive age (18-45 years), and a structured interview schedule was used to collect the data. Data analysis was done using SPSS version 20.0

**Results:** The results showed that the overall knowledge mean score was 21.53 (SD± 3.722), which showed that women of reproductive age (18-45 years) had inadequate knowledge regarding anaemia, out of 200 women, 184 (92.0%) had inadequate knowledge, 16 (8%) had moderately adequate knowledge, and none of them had adequate knowledge. There was significant association between knowledge and socio-economic status of women ( $P = 0.01$ ).

**Conclusion:** The study concluded that study participants had inadequate knowledge and lack of knowledge about anaemia, which may be the main causes for the increasing prevalence of anaemia among them. The study emphasises the urgent need for comprehensive and inclusive public health initiatives to address the widespread lack of anemia awareness among women of reproductive age, ultimately contributing to better maternal and child health in the broader community.

**Keywords:** Anemia; knowledge; women of reproductive age.

## I. INTRODUCTION

Anemia is a major health problem throughout the world, in that iron deficiency anemia is one of the commonest forms of anemia<sup>1</sup>. The prevalence of iron deficiency anemia is higher in developing countries like India, especially affecting toddler, adolescents and women of childbearing age<sup>2,3</sup>.

The manifestations of iron deficiency anemia differ in different age groups, such as increased risk of infectious diseases, affect the cognitive and motor development and mortality in children and adolescents. It will cause premature delivery, low birth weight and also increased risk of maternal mortality rate during pregnancy<sup>3</sup>. Among women in reproductive age the symptoms may not be noticeable in the beginning it is like an iceberg, but paleness, fatigue, low blood pressure can be manifested later. In severe cases, there will be shortness of breath and chest pain, which is an evidence of inadequate perfusion and oxygenation of the major organs<sup>5</sup>. These factors can worsen the health conditions of women such as lung diseases, cardiovascular diseases and heart attack, ultimately death<sup>4,13</sup>. And also result in obstetrical risk such as foetal complication and increased infant and maternal mortality<sup>21</sup>.

In India, two-third of women of childbearing age are estimated to suffer from iron deficiency anemia<sup>9,30</sup>. Reports from National Nutrition Monitoring Bureau 2017 indicate that 15% of all maternal deaths are attributed to anemia. The highest prevalence of anemia among women in India is a burden for them, for their families, and for the economic development



and productivity of the country <sup>21</sup>. Iron deficiency anaemia occurs more often in women than in men; the main reason is excessive loss of iron or demand of iron associated with menstruation and pregnancy <sup>10,28</sup>.

Nearly 400 million women worldwide suffer from iron deficiency anaemia. In India, according to the National Family Health Survey 5, which was undertaken between 2019 and 20, the Prevalence of anaemia among women increased from 53% in NFHS-4 to 57% in NFHS-5. In Karnataka, the incidence of anaemia among married women was about 47.7% among rural married women and 46.7% among urban married women, which seems to be very high <sup>6,7</sup>.

In developing countries like India, where iron deficiency anaemia is more prevalent and high, many women conceive shortly after marriage<sup>5,27</sup>. It is well known that pre-conceptional iron deficiency has adverse effects on pregnancy outcomes. A research study done by the Department of Medicine, St. John's Medical College Hospital, reported that the occurrence of restless leg syndrome was significantly higher among those who had iron deficiency anaemia<sup>7</sup>.

The initial symptoms of iron deficiency anemia in women can go unnoticed. In severe cases, inadequate oxygen supply to major organs can lead to health problems such as kidney failure, lung diseases, and cardiovascular diseases, ultimately leading to death <sup>1,4,8</sup>. In India, iron deficiency anemia is influenced by poverty, illiteracy, and lack of knowledge about dietary requirements. Measures to prevent iron deficiency anemia include consuming iron-rich foods from both animal and plant sources, and fortifying food with iron at the community level<sup>9,15,22</sup>. The present study was undertaken to enhance women's understanding of health-related factors and identify critical influences, ultimately fostering a positive shift in their attitudes toward health.

## II. MATERIAL AND METHODS

**Sample and Sampling Technique:** The present study's sample comprises married women aged 18-45 who were able to understand and speak Kannada and English. Antenatal and postnatal mothers and women who were under treatment for anaemia were excluded from the study. In the present study, 200 women of reproductive age (18-45 years) were selected using a systematic random sampling technique.

**Description of Instrument:** The instrument used in this study consists of two sections, which are as follows. Section I consisted of Demographic variables, including women's age, marital status, number of children, type of family, religion, dietary habits, education, occupation, and family income. Section II consisted of a structured knowledge questionnaire for assessing knowledge regarding anaemia. It consisted of 27 items, each with 3 options, and the total score was 50. The classification of knowledge scoring is given below

**Table 1 Classification of Knowledge Scoring**

SL. NO	Content	Score	Percentage	Total Score
1	Inadequate	<27	<50%	50
2	Moderately adequate	27-36	50-75%	
3	Adequate	>36	>75%	

**Reliability:** The reliability of the tool was established using the split-half method, and coefficient correlation was done with the help of Karl Pearson's formula. The reliability score obtained was  $r=0.8448$  for the knowledge questionnaire, which showed that the knowledge questionnaire was highly reliable. Hence, the tool was considered statistically reliable for the main study.

**Procedure For Data Collection:** Data were collected during June and July 2020 in an urban slum in Bangalore, Karnataka. Using systematic random sampling, every fifth household was selected for the study purpose, and only one eligible, willing respondent from each household was included in the study. Overall, 200 women of reproductive age (18-45 years) from 200 households were interviewed face to face.

**Methods of data analysis:** The data obtained was analyzed on the basis of the objectives of the study using SPSS 20.0 version

## III. RESULT

### Description of demographic variables of women of reproductive age (18-45 years).

For the current research, 200 women of childbearing age (18-49 years old) were chosen. Of the 200 women, the largest group (41.5%) fell within the 18-25 age range, totaling 83 women. 36% (73 women) were between 25-35 years old, and the remaining 22% (44 women) were aged 35-45. The majority, 94.5%, were married (189 women), while 4.5% were widows (9 women) and 1% were divorcees (2 women). Regarding children, 30% (60 women) had more than 2, 28% (56 women) had 2, 25% (50 women) had one child, and the remaining 17% (34 women) had no children. In terms of family structure, 67.5% (135 women) were from nuclear families, and 32.5% (65 women) were from joint families. Religion-wise, 70.5% (141 women) were Hindu, 27.5% (55 women) were Muslim, and 2% (4 women) were Christian. When it came to socioeconomic status, 91% (182 women) were from lower class families, 4.5% (9 women) were from lower middle-class families, and 4% (8 women) were from upper lower class families, while 0.5% (1 woman) was from an upper middle-class family.



**Knowledge level of women of reproductive age (18-49 years) regarding anemia**

**Table 2 Knowledge Score obtained**  
 n=200

Variable	N	Mean	SD
Score	200	21.53	3.742

The results of this study showed that the overall knowledge mean score was 21.53 (SD± 3.722), which showed that women of reproductive age (18-45 years) had inadequate knowledge regarding anemia. Out of 200 women, 184 (92.0%) had inadequate knowledge, 16 (8%) had moderately adequate knowledge, and none had adequate knowledge.

**Table 3 Knowledge level of women of reproductive age (18-45 years) regarding anemia**  
 n=200

Sl no	Content	Max scores	Knowledge Level					
			Inadequate		Moderately adequate		Adequate	
			No	%	No	%	No	%
1	Meaning of anemia	2	34	17	91	45.5	75	37.5
2	Causes of anemia	3	184	92	13	6.5	3	1.5
3	Risk factors	2	17	8.5	170	85	13	6.5
4	Signs and symptoms of anemia	6	165	82.5	34	17	1	0.5
5	Diagnosis of anemia	3	186	93	9	4.5	5	2.5
6	Ill effects of anemia on	8	123	61.5	76	38	1	0.5
7	Treatment of anemia	5	183	91.5	0	0	17	8.5
8	Prevention of anemia	16	127	63.5	73	36.5	0	0
9	Daily requirement of iron	3	91	45.5	95	47.5	14	7
10	Complications	2	47	23.5	143	71.5	10	5

Data reveals that the majority i.e., 184 (92%) women of reproductive age of them had inadequate knowledge regarding the causes of anemia. Majority i.e., 165 (82.5%) of them had inadequate knowledge regarding signs and symptoms of anemia, 183 (91.5%) of them had inadequate knowledge regarding treatment of anemia and 127 (63.5%) had inadequate knowledge regarding prevention of anemia. Only 17 (8.5%) of them had adequate knowledge regarding treatment of anemia

**IV.DISCUSSION**

**Demographic Characteristics and Their Implications**

The study found that among the 200 women, there was a diverse range of ages. The largest group (41.5%) was between 18 and 25 years old, a critical time for a woman’s reproductive health. This suggests that efforts to prevent anemia and other health issues should focus on this age group. Most participants (94.5%) were married, indicating a strong link between marital status and reproductive health needs. 30% of the women had more than two children, which increases the risk of anemia. This shows the importance of providing nutritional

support and education, especially for women with multiple children. Prioritizing these interventions can improve anemia prevention and overall reproductive health outcomes. The study revealed a noteworthy contrast in the educational achievements of the participants. Notably, 31% were found to be illiterate, while only 3% had completed graduate-level education. This disparity is concerning, as literacy plays a critical role in health literacy—the capacity to obtain, comprehend, and utilize information to make well-informed health decisions. The observed association between lower educational attainment and insufficient knowledge about anemia aligns with existing literature underscoring the pivotal role of education in health awareness and outcomes<sup>14,17,25</sup>. The high prevalence of illiteracy among the participants underscores the necessity for health education programs that cater to diverse educational levels, potentially through visual aids, community workshops, and non-text-based educational tools<sup>19,30,34</sup>.



### **Anemia Knowledge Deficits**

The study outlined a noteworthy deficiency in knowledge regarding anemia among the participants. The average knowledge score was 21.53 (SD± 3.722), indicating an overall inadequacy of knowledge about anemia among the women. A substantial 92% of the participants demonstrated a lack of sufficient knowledge about the causes of anemia, while 91.5% were uninformed about effective treatment options. This widespread lack of awareness raises concerns as it directly impacts the participants' ability to prevent and manage anemia, posing significant implications for maternal and child health. Furthermore, 82.5% of the participants displayed inadequate knowledge about the signs and symptoms of anemia, potentially leading to delayed diagnosis and treatment, thereby increasing the risk of severe health outcomes.

Additionally, the study revealed that none of the participants possessed adequate overall knowledge about anemia, highlighting a critical gap in health education. This dearth of knowledge likely contributes to the high prevalence of anemia observed in similar populations. The findings correspond with previous studies reporting low levels of health literacy and awareness about anemia, particularly among women of reproductive age in low-resource settings. This knowledge gap underscores the need for comprehensive educational interventions focusing on improving awareness and understanding of anemia's causes, symptoms, treatment, and prevention<sup>13,18,23</sup>.

### **Association Between Knowledge and Demographic Variables**

The research investigated the relationship between levels of knowledge and specific demographic factors. It was found that family income was significantly linked to knowledge levels ( $P < 0.05$ ). This underscores the influence of socioeconomic elements in determining health knowledge and access to healthcare resources. It was observed that women from higher-income households demonstrated greater knowledge about anemia, likely attributed to improved access to educational resources, healthcare services, and nutritional information. This aligns with global studies that highlight income as a key determinant of health knowledge and outcomes<sup>29,30,32</sup>.

In contrast, other demographic variables such as age, number of children, education, and hemoglobin levels did not display a significant association with anemia knowledge. This suggests that while socioeconomic status plays a crucial role, other variables may not independently predict anemia knowledge to the same extent. Notably, the lack of substantial association between education and anemia knowledge is significant, considering the well-established connection between education and health literacy. This may indicate that factors beyond education, such as the quality of education or access to health information, exert influence on health knowledge even within educational categories<sup>24</sup>.

### **Public Health Implications**

The study highlights the need for tailored educational initiatives to address the gaps in knowledge about anemia among women of childbearing age. It emphasizes the importance of prioritizing public health interventions to raise awareness about anemia, particularly among women with lower income and educational attainment. Additionally, the study underscores the significance of addressing socioeconomic barriers to healthcare access and the necessity for culturally sensitive and readily accessible interventions for effective anemia prevention and management, particularly in underprivileged communities.

### **V. CONCLUSION**

Addressing these knowledge deficits through targeted educational interventions is crucial for improving anaemia awareness and management. Public health strategies must focus on creating accessible, culturally sensitive educational programs that cater to the needs of women across all socioeconomic strata. By empowering women with the necessary knowledge and resources, we can significantly reduce the prevalence of anaemia and enhance the overall health outcomes for women in this population. This study also emphasizes the urgent need for comprehensive and inclusive public health initiatives to address the widespread lack of anaemia awareness among women of reproductive age, ultimately contributing to better maternal and child health in the broader community.

### **VI. REFERENCE**

- [1]. Why women are anemic in India.(2008) available from: URL: <http://www.women.ygoy.com/2008>
- [2]. Robbin and Cortran. (2010) Pathological basis of disease. 7<sup>TH</sup> edition, Elsevier publication. (Pg643-6).
- [3]. Allen. (2000) Anemia and iron deficiency have effects on pregnancy outcomes. The American Journal Of Clinical Nutrition, 1280-84.
- [4]. Beck and Hacker's (2010) .Textbook of gynecologic oncology, 17<sup>th</sup> edition, Lippincott Wilkins publication, (Pg-709)
- [5]. Anemia In Women. available from: URL: <http://www.bolohealth.com>
- [6]. The National Family Health Survey -5 (2019-20) reports. Available from: URL: <http://des.kar-nic.in/ptc/MENWOMEN>
- [7]. D'souza and Rangarajan (2007). Restless leg syndrome in an Indian patient having iron deficiency anemia in a tertiary care hospital. Journal of sleep medicine. (Pg247-251)
- [8]. K V Krishna das. (2011) Text book of medicine, 5<sup>th</sup> edition, Jaypee publications. (Pg 164-165).
- [9]. Park K.(2012). Textbook of preventive and social medicine, 20th edition, M/S Banarsidas Banana Publishers, (Pg. 559).



- [10]. Kothari C R. (2012). Research Methodology Method and techniques. Second Edition, New age international P Ltd publishers, Delhi, (Pg200).
- [11]. Toteja, Singh, Dhillon, Saxena, Ahmed, et, al. (2006). Prevalence of anemia among pregnant women and adolescent girls in 16 districts of India. Food and Nutrition Bulletin. (Pg27-29)
- [12]. Umeta, Haidar, Demisse, Akalu, Ayana (2008). Iron deficiency anemia among women of reproductive age in nine administrative regions of Ethiopia. Ethiopia journal of health. (Pg 252-258).
- [13]. Raghuram V, Manjula anil, Jayaram. (2012). Prevalence of anemia amongst women in the productive age group in a rural area in south India. International journal of biological & medical research. [Online] [Cited (Pg1482-1484).
- [14]. Panigrahi, Bikash Sahoo. (2012). Nutritional anemia and its epidemiological correlates among women of reproductive age in an urban slum of Bhubaneswar, Orissa. (Pg 250-53)
- [15]. Bentley and Griffiths. (2003). The burden of anemia among women in India. European Journal of Clinical Nutrition. (Pg 52–60).
- [16]. Ansari, Ali, Aziz, Ara, Nagina, Tahir. (2009). Nutritional iron deficiency in women of childbearing age- what to do. Journal of Ayub Medical College Abbottabad. (Pg21-3)
- [17]. Sun. (2010). Challenges with Iron Deficiency Anemia among Women of Reproductive Age in Saharawi Refugee Camps, Tindouf, Algeria. Open Digital Archive. (Pg 10-117)
- [18]. Malhotra, Kumari, Kumar and Varma S. (2004). Prevalence anemia in adult rural population of northern India. Journal of association of physicians of India.
- [19]. Pala and Dunder.(2008). Prevalence and risk factors of anemia among women of reproductive age in Bursa, Turkey. Indian journal of medicine, September. (Pg282-286)
- [20]. Thankachan, Muthayya, Walczyk, Kurpad, and Hurrell. (2007) An analysis of the etiology of anemia and iron deficiency in young women of low socioeconomic status in Bangalore, India. Food and Nutrition Bulletin. (Pg28-30)
- [21]. Simon, Peter, Donald. (2008). Hookworm-related anemia among pregnant women: a systematic review. Journal of PLoS Neglected Tropical Diseases. (Pg 291)
- [22]. Joharah M. Al-Quaiz. (2001). Iron deficiency anemia Saudi. Saudi Medical Journal (Pg491-496)
- [23]. Pasricha, Caruana, Phuc, Casey, Jolley, Kingsland. (2008). Anemia, Iron Deficiency, Meat Consumption, and Hookworm Infection in Women of Reproductive Age in Northwest Vietnam. American Journal of Tropical Medicine Hygiene. (Pg375–381)
- [24]. Thankachan, Muthayya, and Hurrell. (2008) An analysis of the aetiology of anaemia and iron deficiency in young women of low socioeconomic status in Bangalore, India. Food and Nutrition Bulletin. (Pg218-22)
- [25]. Rao, Joshi, Dhide, Puranic and Kanade. (2010). Social dimensions related to anemia among women of childbearing age from rural India. Journal of Public Health Nutrition. (Pg 23-28)
- [26]. Ansari, Liaquat ali, Aziz, ara, Nagina. (2009) Nutritional iron deficiency in women of childbearing age, in Karachi, Pakistan. Journal of Ayub Medical College Abbottabad.
- [27]. Sultan. (2007). Anaemia among female college students attending the University of Sharjah: prevalence and classification.
- [28]. D Shojaeizadeh. (2001). Study on Knowledge, Attitude and Practice of Secondary School Girls in Qazvin on Iron Deficiency Anemia. Iranian Journal of Public Health. (Pg53-56)
- [29]. F. Moradi, Mohammadi A. Kadivar and Masoumi. (2007). Knowledge and practice of pregnant women in Fars province about intake of iron supplements. Journal of Acta Medica Iranica.
- [30]. Mirzoyan.(2008). Iron-Deficiency Anemia in Pregnancy: Assessment of Knowledge, Attitudes and Practices of Pregnant Women in Yerevan. (Pg260-7).
- [31]. Upadhyay, Kumar, Singh Raghuvanshi and B. B. Singh. (2001). Nutritional Status and Knowledge of Hill Women on Anemia: Effect of Various Socio-demographic Factors. Journal of Human Ecology (Pg29-34)
- [32]. Kabir, Shahjalal, Saleh, Wahida. (2010). Dietary pattern, nutritional status, anemia and anemia-related knowledge in urban adolescent college girls of Bangladesh. Journal of Pakistan Medical Association. (Pg28-33)
- [33]. Kalimbara AA, and Chilima. (2009). Maternal knowledge and practices related to anemia and iron supplementation in rural Malawi. African Journal Of Food Agriculture Nutrition Development. (Pg 550-564)
- [34]. Hussain and Shu. (2010). Awareness of iron deficiency anaemia among women of reproductive age in Hubei, China. Asian Journal Of Medical Sciences. (Pg12-13).

# IJEAST

INTERNATIONAL JOURNAL  
OF ENGINEERING APPLIED SCIENCE  
AND TECHNOLOGY

## ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

## FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



### PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



### OPEN ACCESS

Free and unrestricted access to research for all.



### GLOBAL REACH

Connecting researchers and professionals worldwide.



### TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website  
[www.ijeast.com](http://www.ijeast.com)



INTERNATIONAL JOURNAL  
OF ENGINEERING APPLIED SCIENCE  
AND TECHNOLOGY

✉ [editor@ijeast.com](mailto:editor@ijeast.com)

🌐 [www.ijeast.com](http://www.ijeast.com)

📍 India



2455-2143