



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 7 ISSUE : 03 Print / Issue Publication Date: 29-Aug-2022



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2022.v07i03.030

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



STUDY ON AMBIENT AIR QUALITY AND EXCEEDANCE FACTOR FOR SOME CITIES OF HARYANA, INDIA

Sadhana Chaurasia

Head, Dept. of Energy & Environment,
MGCGV, Chitrakoot, Satna, M.P. 485334

Swatesh Gupta

PG Student, Dept. of Energy & Environment,
MGCGV, Chitrakoot, Satna, M.P. 485334

Abstract: This Study presents an analysis of the variation of concentration of air pollutants namely PM_{10} , $PM_{2.5}$, SO_x and NO_2 at some city of Haryana, in summer season. The results reveal that gaseous pollutants such as SO_x and NO_2 were within the permissible limits while particulate matter were found higher than permissible NAAQS limit. Highest level of PM_{10} ($198.78 \mu g/m^3$) was found in month of June at Sonipat. It may pose detrimental effect on human and environmental health. In this study AQI and Exceedance factor was also calculated. The AQI for all the stations were found in moderate range (101-200). Exceedance factor for PM_{10} was found in critical range at all the sampling stations and for $PM_{2.5}$ it was found in high pollution range at all stations except Panipat.

Keyword: AQI, Exceedance Factor, NAAQS, Air Pollution.

I. INTRODUCTION:

Air pollution is a serious worldwide environmental problem. After decades of industrialisation, air pollution has become a major environmental issue for both developed and developing countries (Chaurasia et. al. 2020). Poor air quality has chronic effect on environment (Bhuyan et al., 2010). Air pollution seriously damages material sources such as building, various sculptures, and also vegetation. It may be due to particulate matter dispersed in it or gaseous

pollutants completely miscible with it in all proportions. Gaseous pollutants such as SO_2 , NO_x , CO_2 , etc., dispersed in air are the major source of air pollution. (Sarasamma and Narayanan, 2014). Every year large quantities of pollutants are discharged into the environment from ever increasing production of goods and from burning of fossil fuels to generate energy needed to sustain industrial and domestic activities. Awareness of air contamination and measures to monitor and control air quality are inadequate considering the rapidity of increase in pollution levels. (Balashanmugam et al., 2012)

Study Area: Haryana is a state in northern India located at $29.05^\circ N$, $76.08^\circ E$ and approximately 210-275 m.a.s.l. The state has an administrative area of 44,000 km^2 . The population is about 25.4 million. The temperature ranges from $45^\circ C$ to $47^\circ C$ in pre-monsoon and $2^\circ C$ to $5^\circ C$ in winter. The city are dominated by various small, medium and large-scale industries. From 1966 to 1997, the total number of industries has increase up to 7 time, whereas a 75 times increase in total number of vehicles has been registered. Six major cities namely, Manesar, Faridabad, Panipat, Bawal, Hisar and Sonipat were selected for the ambient air quality monitoring (Table: 3-6). Which have observed considerable growth in commercial and industrial sector during last few years.



Figure1Map of Haryana showing sampling stations

Methodology: Ambient air monitoring was conducted during summer season from March 2022 to June 2022.8-hoursampling was done forPM₁₀, PM_{2.5}, SO_x and NO₂.APM-460 respirable dust samplers (RDS) with provision for gaseous sampling APM-415 (Envirotech, New Delhi) was used for measuring the concentrations of PM₁₀, PM_{2.5}, NO₂, and SO_x, in the ambient air. The sampling inlet was placed 1-3 meter above the ground level, depending upon the site available for the RDS. Atmospheric air was drawn for ~8 hours through the cyclone and 20 X 25 cm

glass fiber filter (GFF) sheet at a flow rate of 1.0 to 1.2 m³min and finally the average flow rate was calculated. In the present study, an attempt has been made to assess the prevailing concentration of the PM₁₀, PM_{2.5}, SO_x and NO₂,in the fast-growing urban centers of Haryana(CPCB: 2003). AQI is calculated as per steps given in the CPCB report and computed on Microsoft Excel software and Exceedance factor is used to identify the level of pollution (critical, high, moderate, and low). Following is the equation to find the exceedance factor(Kumar et. al. 2021);

$$\text{Exceedance Factor} = \frac{\text{The annual average concentration of critical pollutant}}{\text{The annual standard for a particular pollutant}}$$

Table1Showing station code and stationname

Station Code	Station Name
S1	Manesar
S2	Faridabad
S3	Panipat
S4	Bawal
S5	Hisar
S6	Sonipat



II. RESULTS AND DISCUSSION:

The sampling and analysis of ambient air quality parameters for six selected station was done in summer season from March 2022 to June 2022. 8 hours sampling was done for all the parameter.

The maximum concentration of PM 2.5 was 71.39 $\mu\text{g}/\text{m}^3$ at S2 and minimum 53.42 $\mu\text{g}/\text{m}^3$ at S5 the concentration of PM2.5 was found in the range of 53.42-71.39 $\mu\text{g}/\text{m}^3$ (Table-3, Figure-2). The average concentration of PM2.5 was found higher than the standard limit (Table-2) except S3.

The maximum concentration of PM₁₀ was 189.41 $\mu\text{g}/\text{m}^3$ at S6 and minimum 151.33 $\mu\text{g}/\text{m}^3$ at S3 the Concentration of PM₁₀ was found in the range of 151.33 – 189.41 $\mu\text{g}/\text{m}^3$

(table-4, Figure-3). The average concentration of PM₁₀ was found higher than the standard limit (Table-2) at all the station.

The maximum concentration of SO_x was 46.25 $\mu\text{g}/\text{m}^3$ at S3 and minimum 35.72 $\mu\text{g}/\text{m}^3$ at S1 the Concentration of SO_x was found in the range of 35.72 – 46.25 $\mu\text{g}/\text{m}^3$ (Table-5, Figure4). The average concentration of SO_x was found within the standard limit (Table-2) at all the station.

The maximum concentration of NO₂ was 75.98 $\mu\text{g}/\text{m}^3$ at S6 and minimum 70.55 $\mu\text{g}/\text{m}^3$ at S5 the Concentration of NO₂ was found in the range of 70.55 – 75.98 $\mu\text{g}/\text{m}^3$ (table-6, Figure-5). The average concentration of PM₁₀ was found within the standard limit (Table-2) at all the station.

Table2 National Ambient Air Quality Standard (2009)

S.NO.	Name of Pollutant	Time weighted average	Concentration in ambient air	
			Industrial, residential, Rural & other Area	Ecologically sensitive area (notified by central government)
1	SO _x ($\mu\text{g}/\text{m}^3$)	Annual	50	20
		24 hours	80	80
2	NO ₂ ($\mu\text{g}/\text{m}^3$)	Annual	40	30
		24 hours	80	80
3	PM ₁₀ ($\mu\text{g}/\text{m}^3$)	Annual	60	60
		24 hours	100	100
4	PM _{2.5} ($\mu\text{g}/\text{m}^3$)	Annual	40	40
		24 hours	60	60

Source: CPCB, 2009

Table 3 Monthly Average Concentration of PM_{2.5}

Month	S1	S2	S3	S4	S5	S6
March	46.59	68.77	44.8	61.96	58.06	66.96
April	54.11	70.82	52.68	69.88	59.08	67.01
May	67.29	72.51	56.89	74.9	60.91	70.34
June	72.86	73.45	59.32	77.62	62.58	71.08
Mean	60.21	71.39	53.42	71.09	60.16	68.85
±SD	12.01	2.06	6.37	6.88	2.00	2.17
Max	72.86	73.45	59.32	77.62	62.58	71.08
Min	46.59	68.77	44.8	61.96	58.06	66.96

Table 4 Monthly Average Concentration of PM₁₀

Month	S1	S2	S3	S4	S5	S6
March	110.41	153.18	123.27	149.65	142.61	174.31
April	149.95	155.93	158.99	153.98	153.98	189.67



May	188.49	159.87	155.63	163.38	163.38	194.89
June	195.05	164.88	167.43	171.63	169.75	198.78
Mean	160.98	158.47	151.33	159.66	157.43	189.41
±SD	39.14	5.08	19.35	9.82	11.81	10.74
Max	195.05	164.88	167.43	171.63	169.75	198.78
Min	110.41	153.18	123.27	149.65	142.61	174.31

Table 5 Monthly Average Concentration of SO_x

Month	S1	S2	S3	S4	S5	S6
March	26.25	39.67	36.98	38.51	31.83	38.9
April	32.78	41.32	48.98	41.94	35.48	33.51
May	40.01	43.44	49.18	44.96	41.13	42.96
June	43.85	44.79	49.86	45.84	43.03	44.74
Mean	35.72	42.31	46.25	42.81	37.87	40.03
±SD	7.81	2.26	6.19	3.32	5.15	4.99
Max	43.85	44.79	49.86	45.84	43.03	44.74
Min	26.25	39.67	36.98	38.51	31.83	38.9

Table 6 Monthly Average Concentration of NO₂

Month	S1	S2	S3	S4	S5	S6
March	67.38	72.66	61.83	70.38	64.01	69.11
April	63.83	73.83	68.28	71.41	68.34	74.88
May	76.93	75.05	75.91	73.62	72.66	77.61
June	79.86	78.66	76.82	75.81	77.18	82.3
Mean	72.00	75.05	70.71	72.81	70.55	75.98
±SD	7.62	2.60	7.05	2.42	5.66	5.51
Max	79.86	78.66	76.82	75.81	77.18	82.3
Min	67.38	72.66	61.83	70.38	64.01	69.11







Table 7 Air Quality Index For various stations

Station code	AQI	Remark	Health Effects
S1	140.5	Moderate	Breathing discomfort to the people with lung
S2	139.75	Moderate	Breathing discomfort to the people with lung
S3	134.25	Moderate	Breathing discomfort to the people with lung
S4	144.5	Moderate	Breathing discomfort to the people with lung
S5	138.25	Moderate	Breathing discomfort to the people with lung
S6	159.75	Moderate	Breathing discomfort to the people with lung




Table 8 Air Quality Index Value Remark and Health Effects

Index Value	Remark	Health Effects
0-50	Good	Minimal Impact
51-100	Satisfactory	Minor breathing discomfort to sensitive people
101-200	Moderate	Breathing discomfort to the people with lung
201-300	Poor	May cause breathing discomfort to people on prolonged exposure and discomfort to people with heart disease
301-400	Very Poor	May cause respiratory illness to the people on prolonged exposure. Effect may be more pronounced in people with lung and heart disease
>401	Severe	May cause respiratory effects even on healthy people and serious health impact on people with lung/heart diseases. The health impact may be experienced even during light physical activity

AQI: The air quality index is used to express the magnitude of air pollution of an area. AQI for various station is given in table 7, figure 6 the range of AQI was found 134.25 - 159.75. All the stations were found in the moderate air quality range (Table -8) it can affect the people suffering for breathing discomfort and lung problems. Highest AQI was found for S6 followed by S4


Exceedance factor: The exceedance factor is the average concentration of critical pollutants and their corresponding national air quality standard. According to their critical pollution level exceedance factor divided into various categories;

Table 9 Exceedance factor with their respective range

Level of pollution	Exceedance factor
Low pollution	< 0.5
Moderate pollution	0.5-0.9
High pollution	1.0-1.4
Critical pollution	>1.5

Table 10 Exceedance Factor for Various Sampling Stations

Station/Parameter	S1	S2	S3	S4	S5	S6
PM2.5	1.00	1.19	0.89	1.18	1.00	1.15
PM10	1.61	1.58	1.51	1.60	1.57	1.89
SOx	0.45	0.53	0.58	0.54	0.47	0.50
NO ₂	0.9	0.94	0.88	0.91	0.88	0.95



Exceedance factor suggests that PM_{2.5} concentration at S3 was found in moderate pollution level and remaining all the station were found in high pollution level. Exceedance factor for PM₁₀ was found in critical level of pollution at all the station. Exceedance factor of SO_x at S1 and S5 was found in low pollution level and remaining all the station were found in moderate pollution level. Exceedance factor of NO₂ was found in moderate to high pollution level (Table-10).

III. CONCLUSION:

This study reveals that the particulates pollutants, PM₁₀ and PM_{2.5} are mostly higher than permissible limits at all the station and SO_x and NO₂ was found within the permissible limit of NAAQS. The mean AQI was found in the range 134.25 – 159.75. AQI shows that all the stations were in moderate quality level. Exceedance factor for PM_{2.5} was found in high pollution level at S1, S2, S4, S5 & S6 and at S3 exceedance factor was found in moderate pollution level. Similarly, exceedance factor of PM₁₀ shows critical pollution level at all the stations (Srinivas and Sateesh, 2015). Generally, exceedance factor of SO_x & NO₂ were found in moderate range for all station.

IV. REFERENCES:

- [1]. Agrawal G., Mohan D., Rahman H., (2021), "Ambient air pollution in selected small cities in India: observed trends and future challenges", IATSS Research, (pp 19-30)
- [2]. Balashanmugam P., Ramanathan A.R., Nehru V.K. (2012), Assessment of ambient air quality in Chidambaram a south Indian town, Journal of Engineering Science and Technology, (pp 292-302)
- [3]. Bhuyan P.K., Samantray P., Rout S.P., (2010), Ambient Air quality status in Choudwar Area of Cuttak District, Inter National Journal of Environmental Science, (pp 343-355)
- [4]. Chaudhary S., Kumar S., Antil R., Yadav S., (2021), "Air quality before and after covid-19 lockdown phases around new Delhi, India", Journal of Health & Pollution, (pp 1-11)
- [5]. Chaurasia S., Singh R., Tiwari A.K., (2020), Air quality of Chitrakoot during covid-19 pandemic lockdown, International Journal of Scientific Development and Research, (pp 391-394)
- [6]. CPCB (2003), Guideline for ambient air quality monitoring, central pollution control board, ministry of environment and forest, govt. of India, National Ambient Air Quality Monitoring Series: NAAQMS/25/2003-04.
- [7]. CPCB (2009), National Ambient air quality standards, central pollution control board, ministry of environment and forest, govt. of India, Notification New Delhi, 18th November
- [8]. CPCB, Parivesh (2001) - Air Pollution and human health, central pollution control board, moef, new Delhi public of ambient air quality in India - A Review, International Journal of Science Technology & Engineering, (pp 237-244)
- [9]. Kaushik S.P., Tyagi A., Tyagi P.K., Tyagi H., (2013), "Air pollution and its Impact on Human health in Panipat City of Haryana, India, International Journal of Advanced Research, (pp 450-457)
- [10]. Kumar P., Kuldeep, Gautam N., (2021), "An assessment of ambient air quality using AQI and exceedance factor for Udaipur City, Rajasthan



- (India)", WEENTECH Proceedings in Energy, (pp 94-106)
- [11]. Matandirotya N.R., (2021), "Research Trends in the field of ambient air quality monitoring and management in South Africa: A Bibliometric Review", Environmental Challenges, (pp 1-8)
- [12]. Salem A.A., Soliman A.A., El-Haty I.A. (2009), Determination Of nitrogen dioxide, sulfur dioxide, ozone, and ammonia in ambient air using the passive sampling method associated with ion chromatographie and potentiometric analyses, Air Qual Atmos Health, (pp 133-145)
- [13]. Sarasamma, J.D., Narayanan, B.K. (2014) Air Quality Assessment in the Surroundings of KMML Industrial Area, Chavara in Kerala, South India. Aerosol and Air Quality Research, (pp 1769-1778.)
- [14]. Srinivas E., Sateesh K., (2015), "Ambient Air Quality monitoring and Possible health Effects due to Air Pollution in Atchutaapuram, Andhra Pradesh, India, International journal of Science Research, (pp 38-41)

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143