



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 4 ISSUE : 09 Print / Issue Publication Date: 10-Mar-2020



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2020.v04i09.011

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



SEMI-IMPLICIT RATIONAL RUNGE-KUTTA METHOD OF SOLVING SECOND ORDER DIFFERENTIAL EQUATION

Eric Augustine Oje

Department of Mathematical Sciences
Taraba State University, Jalingo, Taraba State,
Nigeria

Aggrey Eric Majuk

Department of Computer Graphics and Image
Processing
University of Debrecen, Debrecen, Hungary

Abstract— There is much work done on implicit and explicit rational Runge-Kutta schemes for second order ordinary differential equations. This motivated the need for semi-implicit scheme since no work has been done on it for second order ordinary differential equations. The research was carried out using Taylor's series and Binomial expansion after some related works were reviewed in the literature. The research was completed in which a one-stage semi-implicit rational Runge-Kutta scheme was derived, the convergence and consistency of the scheme was tested and the scheme was convergent and consistent. The adopted scheme was discovered to perform well in terms of approximating the exact solution of a special case of second order ordinary differential equations.

Keywords— ordinary differential equations, Runge-Kutta, semi-implicit scheme

I. INTRODUCTION

An ordinary differential equation (ODE) is an equation of the form:

$$y' = f(x, y) \quad y(x_0) = y_0 \quad a < x < b \quad (1)$$

Arising from mathematical modeling of a large variety of physical problems in nuclear reactions, delay problems and computer aided designs, perturbation problems or dynamical processes in industries, technological fields and economy of some third world countries affected by inflation and other economic depressions. Often, the solution $y(x)$ of the ordinary differential equation (1) possesses components with fast and slow responses and one in which there is pole or discontinuities. The existing methods of solving this ordinary differential equation includes: Runge-Kutta method, Euler's method, implicit backward difference method proposed by Gear in 1971, Hybrid methods by Ademiluyi (1987),

perturbed polynomial process by Lambert and Shaw, extrapolation process by Evans and Fatunla, Rational functions approximation method by Luke et al.

Runge-Kutta schemes are important family of implicit and explicit iterative methods for approximation of solution of ordinary differential equations. Consider the numerical approximation of second order initial value problems of the form:

$$y'' = f(x, y, y'), \quad y(x_0) = y_0, \quad y'(x_0) = y'_0, \quad a \leq x \leq b \quad (1.1)$$

The general s -stage Runge-Kutta scheme for general second order initial value problems of ordinary differential equations of the form (1) as defined by Jain (1984) is:

$$y_{n+1} = y_n + h y'_n + \sum_{r=1}^s c_r k_r \quad (2)$$

And

$$y'_{n+1} = y'_n + \frac{1}{h} \sum_{r=1}^s c'_r k_r \quad (3)$$

Where



$$k_r = \frac{h^2}{2} f \left(x_n + c_i h, y_n + h c_i y'_n + \sum_{j=1}^r a_{ij} k_j, y'_n + \frac{1}{h} \sum_{j=1}^{\infty} b_{ij} k_j \right), \quad i(1)s \quad (4)$$

With $c_i = \sum_{j=1}^i a_{ij} = \frac{1}{2} \sum_{j=1}^i b_{ij}, \quad i(1)r$

Where $c_i, a_{ij}, b_{ij}, c_r, c'_r$ are constants to be determined. The suitable parameter requires extremely lengthy algebraic manipulation, except for

small values of s (Sharp et al, 1992 and Dormand et al, 1987).

In 1982, Hong Yuan Fu proposed a Rational Runge – Kutta scheme for the integration of this kind of ordinary differential equations. Okunbor (1985), Babatola and Ademiluyi (2000) and Bolaji (2005) worked further on this method and proposed schemes that can be classified as explicit, implicit or semi implicit method. Bolarinwa et al (2012) proposed a two-stage semi-implicit rational Runge-Kutta scheme for solving ordinary differential equation of first order.

From the above, the researcher will focus on developing a scheme for semi-implicit rational Runge-Kutta method of solving ordinary differential equation of second order.

II. DERIVATION OF THE SCHEME

The rational form of (2.1) and (2.2) can be defined as

$$y_{n+1} = \frac{y_n + h y'_n + \sum_{i=1}^r w_i k_i}{1 + y_n \sum_{i=1}^r v_i H_i} \quad (2.5)$$

$$y'_{n+1} = \frac{y'_n + \frac{1}{h} \sum_{i=1}^r w'_i k_i}{1 + \frac{1}{h} y'_n \sum_{i=1}^r v'_i H_i} \quad (2.6)$$

where

$$k_i = \frac{h^2}{2} f \left(x_n + c_i h, y_n + h c_i y'_n + \sum_{j=1}^i a_{ij} k_j, y'_n + \frac{1}{h} \sum_{j=1}^i b_{ij} k_j \right), \quad i = 1(1)r \quad (2.7)$$

$$H_i = \frac{h^2}{2} g \left(x_n + d_i h, z_n + h d_i z'_n + \sum_{j=1}^i \alpha_{ij} H_j, z'_n + \frac{1}{h} \sum_{j=1}^i \beta_{ij} H_j \right), \quad i = 1(1)r \quad (2.8)$$

with constraints

$$c_i = \sum_{j=1}^i a_{ij} = \frac{1}{2} \sum_{j=1}^i b_{ij}, \quad i = 1(1)r$$

$$d_i = \sum_{j=1}^i \alpha_{ij} = \frac{1}{2} \sum_{j=1}^i \beta_{ij}, \quad i = 1(1)r$$

in which



$$g(x_n, z_n) = -z_n^2 f(x_n, y_n) \text{ and } z_n = \frac{1}{y_n}$$

The constraint equations are to ensure consistency of the method, h is the step size and the parameters $a_{ij}, b_{ij}, c_i, d_i, \alpha_{ij}, \beta_{ij}$ are constants called the parameters of method.

Following Abhulimen and Uloku(2012), Bolarinwa(2012) procedures:

- i. Obtain the Taylor series expansion of k_i and H_i about the point (x_n, y_n, y'_n) and binomial series expansion of right-hand side of (2.7) and (2.8).
- ii. Insert the Taylor series expansion into (2.7) and (2.8) respectively.
- iii. Compare the final expansion of k_i and H_i about the point (x_n, y_n, y'_n) to

the Taylor series expansion of y_{n+1} and y'_{n+1} about (x_n, y_n, y'_n) in the powers of h .

Normally the number of parameters exceed the number of equations, these parameters are chosen to ensure that (one or more of) the following conditions are satisfied.

- i. Minimum bound of local truncation error exists
- ii. The method has maximized interval of absolute stability
- iii. Minimized computer storage facilities are utilized.

In equations (2.5), (2.6), (2.7) and (2.8), setting $r=1$ we have:

$$y_{n+1} = \frac{y_n + hy' + w_1 k_1}{1 + y_n v_1 H_1} \quad (2.9)$$

$$y'_{n+1} = \frac{y'_n + \frac{1}{h} w'_1 k_1}{1 + \frac{1}{h} y'_n v'_1 H_1} \quad (2.10)$$

where

$$k_i = \frac{h^2}{2} f \left(x_n + c_1 h, y_n + h c_1 y'_n + \sum_{j=1}^1 a_{ij} k_j, y'_n + \frac{1}{h} \sum_{j=1}^1 b_{ij} k_j \right), i = 1 \quad (2.11)$$

$$H_i = \frac{h^2}{2} g \left(x_n + d_1 h, z_n + h d_1 z'_n + \sum_{j=1}^1 \alpha_{ij} H_j, z'_n + \frac{1}{h} \sum_{j=1}^1 \beta_{ij} H_j \right), i = 1 \quad (2.12)$$

with constraints

$$c_1 = a_{11} = \frac{1}{2} b_{11}$$

$$d_1 = \alpha_{11} = \frac{1}{2} \beta_{11}$$



since we are considering the semi-implicit scheme, $b_{ij} = 0$ for $j > i$.

Where $a_{ij}, b_{ij}, c_i, d_i, \alpha_{ij}, \beta_{ij}, w_1, w'_1, v_1, v'_1$ are all constants to be determined.

Now by adopting a binomial expansion (definition 1.6.10) on equations (2.9) gives

$$\begin{aligned} y_{n+1} &= (y_n + hy'_n + w_1 k_1)(1 + y_n v_1 H_1)^{-1} \\ &= (y_n + hy'_n + w_1 k_1)(1 - y_n v_1 H_1) \\ &= y_n + hy'_n - (y_n^2 v_1 + hy_n y'_n v_1) H_1 + (w_1 - y_n v_1 H_1 w_1) k_1 \quad (2.13) \end{aligned}$$

Similarly the binomial expansion of (2.10) gives

$$\begin{aligned} y'_{n+1} &= \left(y'_n + \frac{1}{h} w'_1 k_1\right) \left(1 + \frac{1}{h} y'_n v'_1 H_1\right)^{-1} \\ &= \left(y'_n + \frac{1}{h} w'_1 k_1\right) \left(1 - \frac{1}{h} y'_n v'_1 H_1\right) \\ &= y'_n + \frac{1}{h} w'_1 k_1 - \left(\frac{1}{h} y_n'^2 v'_1 + \frac{1}{h^2} y'_n w'_1 v'_1 k_1\right) H_1 \quad (2.14) \end{aligned}$$

Now, the Taylor's series expansion of y_{n+1} about x_n gives

$$y_{n+1} = y_n + hy'_n + \frac{h^2}{2!} y''_n + \frac{h^3}{3!} y'''_n + \frac{h^4}{4!} y^{iv}_n + \dots$$

and

$$y'_{n+1} = y'_n + hy''_n + \frac{h^2}{2!} y'''_n + \frac{h^3}{3!} y^{iv}_n + \dots$$

where

$$y''_n = f(x_n, y_n, y'_n) = f_n$$

$$y'''_n = f_x + y'_n f_y + f_n f_{y'} = \Delta f_n$$

$$y^{iv}_n = f_{xx} + y_n'^2 f_{yy} + f^2 f_{y'y'} + 2y'_n f_n f_{yy'} + 2f_n f_{xy'} + f_y' \Delta f_n = \Delta^2 f_n + f_y' \Delta f_n + f_n f_y$$

$$\text{since } \Delta = \frac{\partial}{\partial x} + y'_n \frac{\partial}{\partial y} + f_n \frac{\partial}{\partial y'}$$

Using the Taylor's series of the function of three variables (definition 1.6.8) we have from (2.11),



$$\begin{aligned} \frac{2}{h^2} k_1 = f_n &+ \left(c_1 h f_x + (h c_1 y'_n + a_{11} k_1) f_n + \frac{1}{h} b_{11} k_1 f_{y'} \right) \\ &+ \frac{1}{2!} \left[(c_1 h)^2 f_{xx} + 2 c_1 h (h c_1 y'_n + a_{11} k_1) f_{xy} + 2 c_1 h \left(\frac{1}{h} b_{11} k_1 \right) f_{xy'} + (h c_1 y'_n + a_{11} k_1)^2 f_{yy} \right. \\ &\left. + 2 (h c_1 y'_n + a_{11} k_1) \left(\frac{1}{h} b_{11} k_1 \right) f_{yy'} + \left(\frac{1}{h} b_{11} k_1 \right)^2 f_{y'y'} \right] + \dots \end{aligned}$$

simplifying further and re-arranging the equation in powers of h gives

$$\begin{aligned} k_1 = \frac{h}{2} [b_{11} k_1 f_{yy'} + a_{11} b_{11} k_1^2 f_{yy'}] &+ \frac{h^2}{2} \left[f_n + a_{11} k_1 f_y + c_1 b_{11} k_1 f_{xy'} + \frac{1}{2} a_{11}^2 k_1^2 f_{yy} + c_1 y'_n b_{11} k_1 f_{yy'} \right] \\ &+ \frac{h^3}{2} [c_1 f_x + c_1 y'_n f_y + c_1 y'_n f_y + c_1 a_{11} k_1 f_{xy} + c_1 a_{11} y'_n k_1 f_{yy}] \\ &+ \frac{h^4}{4} [c_1^2 f_{xx} + 2 c_1^2 y'_n f_{xy} + c_1^2 y_n'^2 f_{yy}] + 0(h^3) \end{aligned} \quad (2.15)$$

Equation (2.15) is implicit, which cannot be proceeded by successive substitutions, we assume a solution of k_1 which may be expressed as

$$k_1 = h A_1 + h^2 B_1 + h^3 C_1 + h^4 D_1 + O(h^3) \quad (2.16) \text{ in order to}$$

be able to compare and evaluate coefficients of h

Substituting the values of k_1 of (2.16) into (2.15) expand and re-arranging in powers of h gives

$$\begin{aligned} k_1 = \frac{h}{2} [b_{11} (h A_1 + h^2 B_1 + h^3 C_1) f_{y'} + a_{11} b_{11} (h A_1 + h^2 B_1)^2 f_{yy'}] \\ + \frac{h^2}{2} [f_n + a_{11} (h A_1 + h^2 B_1) f_y + c_1 b_{11} (h A_1 + h^2 B_1) f_{xy'} + a_{11}^2 (h A_1)^2 f_{yy} \\ + c_1 y'_n b_{11} (h A_1 + h^2 B_1) f_{yy'}] + \frac{h^3}{2} [c_1 f_x + c_1 y'_n f_y + c_1 a_{11} (h A_1) f_{xy} + c_1 a_{11} y'_n (h A_1) f_{yy}] \\ + \frac{h^4}{4} [c_1^2 f_{xx} + c_1^2 y'_n f_{xy} + c_1^2 y_n'^2 f_{yy}] + 0(h^5) \end{aligned} \quad (2.17)$$

On equating the powers of h from (2.16) and (2.17) gives



$$A_1 = 0$$

$$B_1 = \frac{1}{2}f_n$$

$$C_1 = \frac{1}{2} \left[c_1 f_x + c_1 y'_n f_y + \frac{1}{2} b_{11} f_n f_{y'} \right] = \frac{1}{2} C_1 \Delta f_n$$

$$D_1 = \frac{1}{4} [c_1^2 \Delta^2 f_n + b_{11} \Delta f_n f_{y'} + a_{11} f_n f_y] \quad (2.18)$$

Then

$$k_1 = \frac{h^2}{2} f_n + \frac{h^3}{2} C_1 \Delta f_n + \frac{h^4}{4} [c_1^2 \Delta^2 f_n + b_{11} \Delta f_n f_{y'} + a_{11} f_n f_y] \quad (2.19)$$

In a similar manner

$$H_1 = h^2 M_1 + h^3 N_1 + h^4 R_1 + 0(h^3) \quad (2.20)$$

where

$$M_1 = \frac{1}{2} g_n$$

$$N_1 = \frac{1}{2} d_1 \Delta g_n$$

$$R_1 = \frac{1}{4} [d_1^2 \Delta^2 g_n + \beta_{11} \Delta g_n g_{z'} + \alpha_{11} g_n g_z] \quad (2.21)$$

and also

$$H_1 = \frac{h^2}{2} g_n + \frac{h^3}{2} d_1 \Delta g_n + \frac{h^4}{4} [d_1^2 \Delta^2 g_n + \beta_{11} \Delta g_n g_{z'} + \alpha_{11} g_n g_z] \quad (2.22)$$

Substituting 2.16 and 2.20 into 2.13 and 2.14

$$y_{n+1} = y_n + h y'_n - (y_n^2 v_1 + h y_n y'_n v_1) (h^2 M_1 + h^3 N_1 + h^4 R_1) + (w_1 - y_n v_1 H_1 w_1) (h^2 B_1 + h^3 C_1 + h^4 D_1)$$

Expanding the brackets and re-arranging in powers of h gives

$$y_{n+1} = y_n + h y'_n + h^2 (w_1 B_1 - y_n^2 v_1 M_1) + h^3 (w_1 C_1 - y_n^2 v_1 N_1 - y_n y'_n v_1 M_1) + 0(h^3)$$

also for



$$y'_{n+1} = y'_n + \frac{1}{h} w'_1 (h^2 B_1 + h^3 C_1 + h^4 D_1) - \left(\frac{1}{h} y_n'^2 v'_1 + \frac{1}{h^2} y'_n w'_1 v'_1 (h^2 B_1 + h^3 C_1 + h^4 D_1) \right) (h^2 M_1 + h^3 N_1 + h^4 R_1)$$

Expanding the brackets and re-arranging in powers of h gives

$$y'_{n+1} = y'_n + h(w'_1 B_1 - y_n'^2 v'_1 M_1) + h^2(w'_1 C_1 - y_n'^2 v'_1 N_1 - y'_n w'_1 v'_1 B_1 M_1) + h^3(w'_1 D_1 - y_n'^2 v'_1 R_1 - y'_n w'_1 v'_1 B_1 N_1 - y'_n w'_1 v'_1 C_1 M_1) + 0(h^3)$$

Comparing the corresponding powers in h

$$\begin{aligned} \frac{1}{2} w'_1 f_n - \frac{1}{2} y_n'^2 v'_1 g_n &= \frac{1}{2} f_n \\ \frac{1}{2} w'_1 C_1 \Delta f_n - \frac{1}{2} y_n'^2 v'_1 d_1 \Delta g_n - \frac{1}{2} y'_n y_n' v'_1 g_n &= \frac{1}{6} \Delta f_n \\ \frac{1}{2} w'_1 f_n - \frac{1}{2} y_n'^2 y'_n g_n &= f_n \\ \frac{1}{2} w'_1 \Delta f_n - \frac{1}{2} y_n'^2 v'_1 d_1 \Delta g_n - \frac{1}{2} y'_n w'_1 v'_1 f_n \left(\frac{1}{2} g_n \right) &= \frac{1}{2} \Delta f_n \end{aligned} \quad (2.23)$$

By using equation

$$g_n = -\frac{f_n}{y_n'^2}, \quad g_x = -\frac{f_x}{y_n'^2} g_z = -2\frac{f_n}{y_n'} + f_y, \quad g_z' = -2\frac{f_n}{y_n'} + f_{y'}, \quad z_n' = -\frac{y_n'}{y_n'^2} \quad (2.24)$$

and

$$\Delta g_n = g_n + z_n' g_n + g_n g_z'$$

using 2.24 into 2.23 we get the following simultaneous equations

$$w_1 + v_1 = 1$$

$$w_1 c_1 + v_1 d_1 = \frac{1}{3}$$

$$w'_1 + v'_1 = 2$$

$$w'_1 c_1 + v'_1 d_1 = 1 \quad (2.25)$$

Equation (2.25) has four (4) with six (6) unknowns; there will not be a unique solution, we choose d_1 to be zero in order to obtain a semi-implicit scheme. There will be a family of one-stage scheme.

Choosing the parameters

By solving (2.25) the following values were obtained so the equation 2.25 is satisfied and makes the scheme semi-implicit

$$w_1 = \frac{2}{3}, v_1 = \frac{1}{3}, c_1 = \frac{1}{2}, a_{11} = \frac{1}{2}, b_{11} = 1, d_1 = \alpha_{11} = \beta_{11} = 0,$$

$$w'_1 = 2, v'_1 = 0$$



$$\text{the scheme obtained is } y_{n+1} = \frac{y_n + hy'_n + \frac{2}{3}k_1}{1 + \frac{1}{3}y_n H_1} \quad (2.26)$$

and

$$y'_{n+1} = y'_n + \frac{2}{h}k_1 \quad (2.27)$$

where

$$k_1 = \frac{h^2}{2} f\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hy'_n + \frac{1}{2}k_1, y'_n + \frac{1}{h}k_1\right) \quad (2.27a)$$

$$H_1 = \frac{h^2}{2} g(x_n, z_n, z'_n) \quad (2.27b)$$

The schemes are semi-implicit as required, since k_1 is implicit and H_1 is explicit.

Choosing the parameters

$$\begin{aligned} w_1 &= \frac{1}{3}, v_1 = \frac{2}{3}, c_1 = a_{11} = \frac{1}{2}, b_{11} = 1, \text{ since } c_1 = a_{11} = \frac{1}{2}b_{11}, w'_1 = 2, v'_1 = 0, d_1 = \alpha_{11} \\ &= \frac{1}{4}, \text{ but } d_1 = \alpha_{11} = \frac{1}{2}\beta_{11} \Rightarrow \beta_{11} = \frac{1}{2} \end{aligned}$$

By solving (2.25) the following values were obtained so the equation 2.25 is satisfied and makes the scheme semi-implicit

The following scheme is obtained

$$y_{n+1} = \frac{y_n + hy'_n + \frac{1}{3}k_1}{1 + \frac{2}{3}y_n H_1}$$

and

$$y'_{n+1} = y'_n + \frac{2}{h}k_1$$

where

$$k_1 = \frac{h^2}{2} f\left(x_n + \frac{1}{2}h, y_n + \frac{h}{2}y'_n + \frac{1}{2}k_1, y'_n + \frac{1}{h}k_1\right)$$

$$H_1 = \frac{h^2}{2} g\left(x_n + \frac{1}{4}h, z_n + \frac{h}{4}z'_n + \frac{1}{4}H_1, z'_n + \frac{1}{2h}H_1\right)$$



III. NUMERICAL PROBLEMS

Example 1.

Solve $y'' + 2y = 0, y(0) = 1, y'(0) = 0$,

The exact solution is given by

$$y(x) = \cos x\sqrt{2}, \quad y'(x) = -\sqrt{2} \sin x\sqrt{2}$$

Example 2.

Consider the equation $y' = (1 + x^2)y, y(0) = 1, y'(0) = 0, x \in [0, 0.1]$

The exact solution is

$$y(x) = e^{\frac{x^2}{2}}$$

IV. DISCUSSION OF RESULTS

The first example 1 is an initial valued second order differential equation. This equation suits the scheme (3.26) as it is a second order ordinary differential equation of the form $f(y)$ that is it is an autonomous system. The results get better with decrease in step size (h) as shown in table 1 in appendix A. 24

The table 2 in appendix A is the result obtained by applying the schemes (3.26) to example 2 evaluated from step size $h=0.01$, as it is a special case of second order ordinary differential equation of the form $f(x,y)$. The result shows that the scheme performed well as the approximate solution approaches the exact solution better as the step size reduces to $h=0.001$ this implies the smaller the value of the step size, the better the approximated solution. This scheme approximates the special second order differential equation better as it is shown in table 2 in appendix A.

V. CONCLUSION

The semi-implicit rational Runge-Kutta scheme obtained in this research performed well in approximating the exact solutions and the result of the approximated solution gets better as the step size tends to zero, as it is shown in the result from table 2 in appendix A. The scheme is convergent as well as it is also consistent. This result implies that the scheme does better approximation on the special second order differential equation.

VI. REFERENCES

1. Abbulimen, C.E. and Uloko, J.A (2012). *A Class of an Implicit Stage-two Rational Runge-Kutta Method for Solution of Ordinary Differential Equations*. Journal of Applied Mathematics and Bioinformatics. 2(3): 17-31
2. Ademiluyi, R.A. (1987) "A Class of Hybrid Methods for the Solution of Ordinary Differential Equations". M.sc thesis, University of Benin, Benin city; Nigeria. Unpublished.
3. Ademuluyi, R.A., Babatola, P.O. (2000). "Implicit Rational Runge-Kutta Scheme for Integration of Stiff ODEs". Nigeria Mathematical Society, (NMS) Journal.
4. Bolarinwa, B., Ademiluyi R. A., Oluwagunwa A. P. and Awomuse B. O. (2012). "A Class of Two-Stage Semi-Implicit Rational Runge - Kutta Scheme for Solving Ordinary Differential Equations". Canadian Journal of Science and Engineering Mathematics, 3(3):99 – 111.
5. Bolarinwa, B. (2005). "A Class of Semi-Implicit Rational Runge-Kutta Scheme for Solving Ordinary Differential Equations with Derivative Discontinuities"; M. Tech Thesis, Federal University of Technology, Akure; Nigeria. Unpublished.
6. Dass, H.K. (2012). *Advance Engineering Mathematics*. Third edition. S. Chand and Company Limited. New Delhi, India
7. Dormand, J.R., El-Milkawy, M.E.A, and Prince, P.J.(1987) "Families of Runge-Kutta-Nystrom formula", IMA Journal of Numerical Analysis: 7: 235-250
8. Dresner, L. (1999), *Applications of Lie's Theory of Ordinary and Partial Differential Equations*, Bristol and Philadelphia: Institute of Physics Publishing.
9. Fudziah, I. (2003). *Embedded Singly Diagonal Implicit Runge-Kutta-Nystrom Method Order 5(4) for the Integration of Special Second Order Ordinary* 27
10. Obimba, D.A. (1993). *Basic Linear Differential Equations*. First Edition. Oganiru Publishing Company, Owerri, Nigeria.
11. Odekunle, M.R. (2001). "Some Semi-implicit Rational Runge-Kutta Schemes". Bagale Journal of Pure and Applied Sciences, 1(1):11-14



12. Okunbor, D.I. (1985); “Runge-Kutta Schemes for Stiff ODEs” M.sc Thesis, University of Benin, Nigeria.
13. Okunuga, S.A., Sofoluwa, A.B., Ehigie, J.O. and Akambi, M.A. (2012). *Fifth Order Two-Stage Explicit Runge-Kutta-Nystrom Method for Direct Integration* 28
14. Fudziah, I. (2009). *Sixth Order Singly Diagonally Implicit Runge-Kutta Nystrom Method With Explicit First Stage for Solving Second Order Ordinary Differential Equations*. European Journal of Scientific Research. 26(4):470-479.
15. Gear, C.W. (1971). “DIFSUB for Solution of ODEs” Math comp. vol. 4 page 185-190.
Graham, R., Knuth, D., Patashnik, O. (1994), “(5) Binomial Coefficients” Concrete Mathematics (2nd ed.) Addison Wesley.
16. Greenberg, M. (1998). Advanced Engineering Mathematics (2nd ed.), Prentice Hall.
17. Hong, Y.F. (1982): “A Class Of A-Stage Explicit Scheme, Computational and Asymptotic Method for Boundary and Interior Layer”. Proceeding of ball 11 Conference, Trinity College Dublin, 236-241.
18. Jain, M.K. (1984). “Numerical Solution of Differential Equations, 2nd Edition”. Wiley Estem Limited. New Delhi, India.
19. Lambert, J.D. (1973). Computational Methods in Ordinary Differential Equations. New York: John Wiley and sons.
20. Oye, N.D., Adey, S.O., Odekunle, M.R., (2004). “A Class of Inverse Runge-Kutta Schemes for the Numerical Integration of Singular Problems”. Journal of Applied Mathematics and Computing. Elsevier: 158: 149-158
21. Raisinghania, M.D. (2012). Ordinary and Partial Differential Equation. Fourth edition. New Delhi, India.
22. Senu, N., Sulaiman, M., Ismail, F. and Othman, M. (2011). *A Singly Diagonally Implicit Runge-Kutta-Nystrom Method for Solving Oscillatory Problems*. International Journal of Applied Mathematics. 4(1);
23. Sharp, P.W. and Fine, J.M. (1992). “Nystrom Pairs for the General Second Order Initial-Value Problems”, Journal of Computational and Applied Mathematics 42: 279-291
24. Stroud, K.A. and Dexter, J.B. (2007). Advance Engineering Mathematics, Sixth Edition. Palgrave Macmillan Company, New York.
25. Usman, A.S., Odekunle, M.R. and Ahmad, M.M. (2013). “A Class of Three Stage Implicit Rational Runge-Kutta Schemes for Approximation of Second Order Ordinary Differential Equations”. The International Institute for Science, Technology and Education. 3(11), 121-130.



APPENDIX A

Table 1: Solution of Example 1 at $h=0.001$

	Exact solutions		Approximated solutions		Error	
H	$y(x)$	$y'(x)$	$y(x)$	$y'(x)$	y	y'
0.001	0.99001665	-0.19933400	1.00000000	-0.00199999	9.98E-03	1.97E-01
0.002	0.96026596	-0.39468800	1.00000000	-0.00399999	3.97E-02	3.91E-01
0.003	0.91134193	-0.58216130	1.00000000	-0.00599997	8.87E-02	5.76E-01
0.004	0.84422141	-0.75801080	1.00000000	-0.00799997	1.56E-01	7.5E-01
0.005	0.76024460	-0.91872540	1.00000000	-0.00998752	2.40E-01	9.09E-01
0.006	0.66108821	-1.06109600	1.00000000	-0.01199998	3.39E-01	1.05E00
0.007	0.54873208	-1.18228010	1.00000000	-0.01399966	4.51E-01	1.17E00
0.008	0.42541959	-1.27985800	1.00000000	-0.01599949	5.75E-01	1.26E00
0.009	0.29361288	-1.35188130	1.00000000	-0.01799927	7.06E-01	1.33E00
0.010	0.99001665	-0.19933400	1.00000000	-0.01999999	9.98E-03	1.79E-01
0.020	0.96026596	-0.39468800	1.00000000	-0.03999200	3.97E-02	3.55E-01
0.030	0.91134193	-0.58216130	1.00000000	-0.05997302	8.87E-02	5.22E-01
0.040	0.84422141	-0.75801080	1.00000010	-0.07993605	1.56E-01	6.78E-01
0.050	0.76024460	-0.91872540	1.00000020	-0.09987515	2.40E-01	8.19E-01
0.060	0.66108821	-1.06109600	1.00000040	-0.11978440	3.39E-01	9.41E-01
0.070	0.54873208	-1.18228010	1.00000080	-0.13965780	4.51E-01	1.04E00
0.080	0.42541959	-1.27985800	1.00000140	-0.15948960	5.75E-01	1.12E00
0.090	0.29361288	-1.35188130	1.00000220	-0.17927390	7.06E-01	1.17E00
0.100	0.15594369	-1.39691200	1.00000330	-0.19900490	8.44E-01	1.20E00
0.200	0.96026596	-0.39468800	1.00053700	-0.39215690	4.02E-02	2.53E-03
0.300	0.91134193	-0.58216130	1.00274900	-0.57416270	9.14E-02	7.80E-03
0.400	0.84422141	-0.75801080	1.00884500	-0.74074070	1.65E-01	1.73E-02
0.500	0.76024460	-0.91872540	1.02222200	-0.88888890	2.62E-01	2.98E-02
0.600	0.66108821	-1.06109600	1.04817100	-1.01694900	3.87E-01	4.41E-02
0.700	0.54873208	-1.18228010	1.09547100	-1.12449800	5.47E-01	5.78E-02
0.800	0.42541959	-1.27985800	1.18040900	-1.21212100	7.55E-01	6.77E-02
0.900	0.29361288	-1.35188130	1.33838800	-1.28113900	1.04E00	7.07E-02
1.000	0.15594369	-1.39691200	1.66666700	-1.33333300	1.51E00	6.36E-02



Table 2: Solution to Example 2 at $h=0.001$

	Exact Solutions		Approximated Solutions		Error	
H	y(x)	y'(x)	y(x)	y'(x)	y	y'
0.001	1.00000050	0.00100000	1.00000000	0.00100000	5.00E-07	0.00E00
0.002	1.00000200	0.00200000	1.00000100	0.00200000	1.00E-06	0.00E00
0.003	1.00000450	0.00300001	1.00000100	0.00300000	3.50E-06	1.00E-09
0.004	1.00000800	0.00400003	1.00000300	0.00400003	5.00E-06	0.00E00
0.005	1.00001250	0.00500006	1.00000400	0.00500006	8.50E-06	0.00E00
0.006	1.00001800	0.00600011	1.00000600	0.00600011	1.20E-05	0.00E00
0.007	1.00002450	0.00700017	1.00000800	0.00700017	1.65E-05	0.00E00
0.008	1.00003200	0.00800026	1.00001100	0.00800026	2.10E-05	0.00E00
0.009	1.00004050	0.00900037	1.00001400	0.00900037	2.65E-05	0.00E00
0.010	1.00000050	0.00100000	1.00001700	0.01000050	1.65E-05	9.00E-03
0.020	1.00000200	0.00200000	1.00006700	0.02000400	6.50E-05	1.80E-02
0.030	1.00000450	0.00300001	1.00015000	0.03001351	1.46E-04	2.70E-02
0.040	1.00000800	0.00400003	1.00026700	0.04003202	2.60E-04	3.60E-02
0.050	1.00001250	0.00500006	1.00041800	0.05006256	4.06E-04	4.51E-02
0.060	1.00001800	0.00600011	1.00060200	0.06010815	5.84E-04	5.41E-02
0.070	1.00002450	0.00700017	1.00082000	0.07017181	7.96E-04	6.32E-02
0.080	1.00003200	0.00800026	1.00107200	0.08025661	1.04E-03	7.23E-02
0.090	1.00004050	0.00900037	1.00135900	0.09036561	1.32E-03	8.14E-02
0.100	1.00005000	0.01000050	1.00501900	0.10050190	4.97E-03	9.05E-02

IJEAST

INTERNATIONAL JOURNAL OF ENGINEERING APPLIED SCIENCE AND TECHNOLOGY



AUTHORS

 Eric Augustine Oje

 Aggrey Eric Majuk



ABOUT IJEAST

International journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high - quality research paper in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

JOURNAL METRICS



H-INDEX

33



i10-INDEX

154



IJEAST
CITATIONS

9,647

(IJEAST CITATIONS)



PEER-REVIEWED
& OPEN ACCESS

PUBLISHED
MONTHLY



PEER REVIEWED

All submission are rigorously peer reviewed to ensure quality.



AUTHORS ARE OUR PRIORITY

We value our authors and recognize their contribution to the advancement of research and knowledge.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



editor@ijeast.com



www.ijeast.com



India



2455-2143