



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 6 ISSUE : 10 Print / Issue Publication Date: 21-Apr-2022



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2022.v06i10.036

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



ENHANCEMENT THE SEEDLING QUALITY OF OKRA THROUGH PRIMING WITH H₂O₂

Md. Rahim Badsha
Ex MS student
Dept. of Horticulture,
Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur.

Fahadul Haque, Mahbub Iqbal
Assistant Director
Bangladesh Agricultural Development Corporation (BADC), Motijheel, Dhaka

Shetul Muna Parvin
Post graduate student
Department of Horticulture Bangladesh Agricultural University
Mymensingh-2202, Bangladesh

Md. Abu Zafur Al Munsur
Additional Deputy Director
Department of Agricultural Extension, Khamarbari, Dhaka 1215

T.M.T Iqbal
Professor
Dept. of Horticulture
Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur.

ABSTRACT - The experiment was conducted to evaluate the physiological fluctuations in the seeds of okra seeds primed with the aqueous solutions of H₂O₂ and to find the suitable strength(s) of H₂O₂ as the priming material for their highest seedling quality. The experiment had eight treatments: T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂: T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%). The seeds were soaked for six hours in those seven media (T₂ to T₈). The seeds were dibbled in wooden seed flats filled-in with coarse sand. The experiment was laid-out in the RCBD with three replications. Data collected for the 13 traits were: % germination, % abnormal seedlings, shoot length, root length, seedling length, shoot dry matter, root dry matter, seedling dry matter, number of secondary roots (>1cm) per seedling, number of true leaves per seedling, relative growth rate, seedling vigor index, and root: shoot ratio (dry weight basis). Except the first two traits, the rest 11 were collected at three stages: 10, 20 and 30 days after dibbling (DAD). It was lucid that H₂O₂ was significantly (P≤0.05%) effective to improve most of the traits noted. But 1% concentration was utmost helpful for both okra and others were toxic

and hindered the maximum parameters for all the three species. Nevertheless, further studies with different varieties of those three vegetables species with variable doses of H₂O₂, priming time and temperature could be explored before drawing valid conclusions.

Key words: Enhancement, Seedling Quality, Okra, Priming, H₂O₂

I. INTRODUCTION

Okra (*Abelmoschus esculentus* L.) of family Malvaceae is an important vegetable crop grown in many tropical, subtropical and warmer temperate areas [1]. Okra is an erect, coarse, robust annual herb in which flowers are borne singly in the leaf axile on peduncles. The plant has malvaceous floral organization with 8-10 very narrow, hairy, bracteoles forming an epicalyx. The leaves are, leathery or rough, large, alternate, cordate divided into 3-7 lobes with notched or toothed margins. Flower borne singly in the leaf axils on peduncles 2 to 5 cm long with malvaceous floral organisation The fruits are light green or sometimes red in colour, long (10-30 cm), beaked, ridged;

more or less oblong hairy capsules that dehiscing longitudinally [2-6].

Seed germination is one of the vital stages in the life cycle of seeded plants. Germination is a very complex process starting with the imbibition of H_2O and involves events related with the transition of a dry quiescent and / or dormant seed to the metabolically active state [7-8]. The emergence of the embryonic axis through structures surrounding the embryo is the final stage of germination [9]. In this link, seed priming is used as a means to enhance seed performance, notably in terms to the rate and the uniformity of germination [10]. Seed priming is known as the pre-sowing approach to govern seed germination and seedling development by modulating pre-germination metabolic activities prior to emergence of the radicle and usually enhances germination rate and plant growth [11]. Various physiological and bio-chemical changes happen in seeds during priming as a result of osmotic conditioning. A wide range of pre-sowing hydration techniques is used to enhance seed germination responses. These include equilibrium under conditions of high humidity [12], soaking in plain H_2O [13] or osmotic solutions [14] and having equilibrium with a matric potential controlling surface [15]. Hydro-priming, osmo-priming (with mannitol or PEG 6000) and halo-priming (with KCl, KNO_3 or calcium salts) are effective for seedling establishment under harsh conditions [16]. The priming enhances rapid and uniform emergence, high vigor and better yield, which has practical utilities, preferably under water stress conditions [17].

Priming of seeds with H_2O_2 leads to break primary dormancy [18]; secondary dormancy provoked by salinity [19] and germination inhibitors [20]. It acts as a stress signal in plants and hence exogenous uses of H_2O_2 in the right dose ameliorates seed germination, reduces time to germinate and seedling growth in many crops [21]. So, it has special roles, especially in invigorating seeds with low vigor including long-term stored seeds in gene banks. So, instead of H_2O , seed priming with aqueous solution of H_2O_2 with the appropriate dose could be an applicable technology [22]. But, information in this connection is lacking with the seeds of the okra. The present study was conducted to evaluate the physiological changes of the seedlings of okra primed with the aqueous solution of eight different concentrations of H_2O_2 and find the best one.

II. MATERIALS AND METHODS

The present research work was conducted at the roof top of Sunway Dormitory near the Bus Terminal, Dinajpur, Bangladesh. The seeds of okra collected from the Popular Seed Limited, Bangladesh: okra cv. Arka, was used as the testing seed materials.

2.1 Experimental treatments

This single factor experiment was designed with eight treatments viz. T_1 (control/no priming), T_2 (hydro-priming in plain H_2O) and other six aquatic solutions of H_2O_2 : T_3 (0.5%), T_4 (1.0%), T_5 (1.5%), T_6 (2.0%), T_7 (2.5%) and T_8 (3.0%). This single factor experiment was set in the Randomized Complete Block Design (RCBD) with three replications.

2.2 Seed flats and their arrangements

The seed were debbled in wooden seed flats. The size of each flat was $50 \times 50 \times 15$ cm. Firstly; the flats were set on the roof top. Then blue polyethylene sheet was spread at the bottom of the flats to protect washing away of sand from the seed flats. Then flats were filled-in with coarse sand.

2.3 Preparation of the required H_2O_2 solutions

The required solutions were prepared by diluting the required amounts of the H_2O_2 (30% strength, Plate 1) with H_2O to get these six concentrations: 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0%.



Plate 1. The container of the H_2O_2 (30%) used in the study

2.4 Priming process

At first, only H_2O and those six solutions were taken in plastic glasses separately. The glasses were marked about the treatments and replications with a permanent glass marker. Then the 200 seeds for each replication were taken in plastic glasses to soak in the desired solutions for six hours in the Laboratory of Horticulture (Plate 2).



Plate 2. Priming process with H_2O_2

2.5 Dibbling the seeds and caring the seedlings

The unprimed and the primed seeds were then dibbling immediately in the seed flats (Plate 3) at the depth of 2cm in lines on the 10th March, 2019 at the distance of 5cm between rows and seeds too. After sowing, the seeds were covered with hyaline polyethylene sheet and concrete poles (at one foot high) to protect the seeds and seedlings from heavy rainfall. Light watering with a watering cane was done as needed. Hand weeding was also done as per need.



Plate 3. Growing the seedlings in the seed flats

2.6 Data collection

The data were collected for % germination and % abnormal seedlings at 10 days of dibbling and shoot length, root length, seedling length, shoot dry matter, root dry matter, seedling dry matter, number of secondary roots (>1cm) per seedling, number of true leaves per seedling, relative growth rate, seedling vigor index, and root: shoot ratio (dry weight basis) at 10, 20 and 30 DAD.

Germination (%) and normal seedling (%) were observed and counted as per the ISTA (2010) rules daily up to 10 DAD. For dry matter the normal seedlings were cut and divided into roots and shoots with a razor blade from each treatment and replication wise. Then those were first sundried separately for two days. After that, those were dried at 80°C for 48 hours in an electric oven (Memmert, ULP 400). Then the dry weights of shoots were recorded up

to four decimal places with an electric digital balance (Ohaus, pioneer pro PA214). Finally, the dry weights were expressed in gram per 100-seedling basis. Those processes were repeated with the normal seedlings only obtained from the 10, 20 and 30 DAD. Relative growth rate (RGR) was calculated as per Williams (1946) formula and Seedling vigor index (SVI) was calculated as per Orchard (1977) and Baki and Anderson (1973) viz. $SVI = \text{Mean seedling (root + shoot) length (MSL) in cm} \times \% \text{ germination (PG)}$

2.7 Statistical analyses

The analyses of variances (ANOVA) were done and the means were separated using Duncan's Multiple Range Test (DMRT). The MSTAT-C Statistical Package program was used for it.

III. RESULTS AND DISCUSSION

3.1 Germination %

The priming treatments had a significant effect ($P \leq 0.05$) on the % germination of okra seeds (Fig. 1). The highest (88.00) and the lowest (79.33%) rates of germination were related to primed seeds with 1% (T_4) and 3% (T_8) H_2O_2 solutions, respectively. The results clearly show that more germination advantage of okra seeds were achieved by priming with H_2O_2 as priming changed the germination mechanism due to improved breaking primary and secondary dormancy and also prevents suffocation. In that case, the highest germination % occurred when the seeds were primed with the 1% H_2O_2 solution (T_4). However, the higher doses (>1%) of H_2O_2 probably caused harmful effects on the germination and so, the germination % became low, even than the unprimed and the hydro-primed seeds. The H_2O_2 levels >2% resulted in severe injury to germinating seeds of bitter melon [23] (John and Duval, 2000). [24] Nandi et al. (2017) found better germination in chilli seeds at the 2% concentration of H_2O_2 . [25] Mustafa (2017) studied cucumber, swamp cabbage, radish and Indian spinach seeds and had better germination at 1% H_2O_2 . While working with bitter melon and bottle melon, [26] Lima (2017) found better germination at 1.5% H_2O_2 . But Iqbal et al. (2001) [27] found higher germination in okra seeds at 2% concentration with H_2O_2 . Furthermore, Kaya et al. (2006) [28] reported that priming increased % germination of sunflower seeds under drought stress. Barba-Espinet et al. (2010) [29] claimed that H_2O_2 increased the germination % of pea seedlings in a concentration-dependent manner.

3.2 Percent abnormal seedlings

There was significant difference ($P \leq 0.05$) for the effects of H_2O_2 concentrations among the treatments in respect of % abnormal seedlings (Fig. 2). The maximum % abnormal seedlings (5.66) were recorded in T_8 while the lowest (2.00%) was counted in T_4 . The results show that the incidences of abnormal seedlings increased significantly



with the rise in H₂O₂ concentrations. In bitter gourd and bottle gourd seeds, [26] Lima (2017) noted the Minimum% abnormal seedlings with 1.5% while the maximum at the 3% H₂O₂. Parallely, [25] Mustafa (2017) recorded the minimum % abnormal seedlings with swamp cabbage, radish and Indian spinach seeds at 1% but the maximum abnormal seedlings with 3% H₂O₂. In addition, Kaya et al. (2006) [28] reported that priming reduced the number of abnormal seedlings of sunflower under drought stress.

T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%)

3.3 Shoot, root and seedling length (cm)

The shoot, root and seedling length were significantly different (P≤0.05) for the effects of H₂O₂ concentrations at all the three DADs (Table 1). At the 10, 20 and 30 DAD, the longest shoot was observed in T₄ (5.99, 7.38 and 9.15 cm, respectively) while the shortest one was in T₈ (5.37, 6.52 and 7.98 cm, respectively). The highest root length (4.91, 13.96 and 16.64 cm) was obtained from T₄ while the lowest (3.53, 11.28 and 15.01cm) was recorded from T₈ at the 10, 20 and 30 DAD, respectively. At the 10, 20 and 30 DAD, the tallest seedling (10.89, 21.39 and 25.80 cm) was measured in T₄ while the shortest one (8.91, 17.81 and 22.99 cm) was recorded in T₈. H₂O₂ influenced the vegetative growth of the seedlings and the ultimate results were the longest seedlings compared to the control one. During the seedling growth, the highest shoot length was found when was primed with 1% H₂O₂ and the lowest was when 3% H₂O₂ solution was used. In bitter gourd and bottle gourd seeds, [26] Lima (2017) measured the minimum shoot and seedling length with 1.5% while the maximum at the 3% H₂O₂. Mostafa (2017) also noted the minimum shoot length with swamp cabbage, radish and Indian spinach seeds at 1% but the maximum shoot and seedling length with 3% H₂O₂. But 1.0% was very helpful for country bean and yard long bean while 1.5% for bottle gourd and bitter gourd seeds. Above 1.5%, others were toxic [26] (Lima, 2017). While working with seeds of two varieties of okra, Iqbal et al. (2001) [27] also achieved nearly similar results with the concentration i.e. 2% H₂O₂. The H₂O₂ treatment, which increased shoot lengths, may be useful in crop production as rapid growth of seedling is vital for the better establishment of seedlings [29] (Ogiwara and Terashima, 2001). Again, the higher doses (>1%) of H₂O₂ probably caused poison effects on shoot length and so, the shoot length became low, even that the unprimed and the hydro-primed seeds. However, the applications of H₂O₂ at doses of 1.8mM each eight days broccoli seedlings, increased the stem length and fresh weight, whereas the dose 1.4mM increased the biomass of broccoli seedlings [30] (Leon-Vargas et al., 2016). The influences of seed priming in improving the root length had also been well documented in ground nut by [31] Rahman et al. (1997). [24] Nandi et al. (2017) noted the maximum root length in seedlings of chilli raised from the seeds treated with the 1.0% H₂O₂.




Fig. 1. Effect of priming seed with H₂O₂ on the % germination of okra

T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%)




Fig. 2. Effect of priming seeds with H₂O₂ on the % abnormal seedlings of okra



Table 1. Effect of priming seed with H_2O_2 on shoot, root and seedling length in okra

| Treatments | Shoot length (cm) | | | Root length (cm) | | | Seedling length (cm) | | |
|----------------|-------------------|--------|---------|------------------|---------|----------|----------------------|--------|----------|
| | 10 DAD | 20 DAD | 30 DAD | 10 DAD | 20 DAD | 30 DAD | 10 DAD | 20 DAD | 30 DAD |
| T ₁ | 5.44b | 6.79bc | 8.59b | 4.35ab | 12.14bc | 15.43bc | 9.76bc | 18.93c | 24.03b-d |
| T ₂ | 5.52b | 7.11ab | 8.44bc | 3.92bc | 12.01bc | 15.03c | 9.44cd | 19.12c | 23.48b-d |
| T ₃ | 5.48b | 6.92bc | 8.48bc | 3.90bc | 12.39b | 15.76a-c | 9.38cd | 19.34c | 24.24bc |
| T ₄ | 5.99a | 7.38a | 9.15 a | 4.91a | 13.96a | 16.64a | 10.89a | 21.39a | 25.80a |
| T ₅ | 5.71ab | 7.12ab | 8.08cd | 3.79bc | 13.40a | 15.18c | 9.50c | 20.48b | 23.26cd |
| T ₆ | 5.47b | 6.86bc | 8.36b-d | 4.63a | 11.84bc | 16.21ab | 10.11b | 18.71c | 24.58b |
| T ₇ | 5.50b | 6.88bc | 8.39b-d | 3.78bc | 12.15bc | 14.97c | 9.28cd | 19.03c | 23.36cd |
| T ₈ | 5.37b | 6.53c | 7.98d | 3.53c | 11.28c | 15.01c | 8.91d | 17.81d | 22.99d |
| LSD (0.05) | 0.313 | 0.379 | 0.395 | 0.580 | 0.886 | 0.901 | 0.5015 | 0.7968 | 1.074 |
| CV % | 3.22 | 3.10 | 2.68 | 8.09 | 4.08 | 3.31 | 2.96 | 2.35 | 2.56 |

The figures with different letters differ among themselves at the 5% level of probability.

T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%)

3.4 Shoot, root and seedling dry matter

There was significant variation ($P < 0.05$) in shoot, root and seedling dry matter (g per 100-seedling basis) accumulation among the treatments at all the three DADs (Table 2). At 10, 20 and 30 DAD, the shoot gathered the utmost amount of dry matter (6.93, 9.16 and 19.64 g, respectively) in T₄ although the minimum (5.74, 6.75 and 15.99 g, respectively) was observed in T₈. The highest dry matter in roots was deposited in T₄ (1.14, 1.64 and 4.46 g) and the minimum dry matter of roots was weighed in T₈ (0.64, 1.15 and 3.63 g) at 10, 20 and 30 DAD, respectively. At the 10, 20 and 30 DAD, the highest dry matter formation in

seedlings was obtained at T₄ (8.08, 10.82 and 24.11 g, respectively) while the least was in T₈ (6.36, 7.90 and 19.62 g, respectively). While dealing with the seeds of bitter melon and bottle gourd, [26] Lima (2017) noted the highest shoot, root and seedling dry matter at 1.5% H₂O₂. [25] Mustafa (2017) recorded the maximum dry matter in shoot root and seedlings of swamp cabbage, radish and Indian spinach primed with 1% H₂O₂ and above 1.5%, others were toxic. Soaking seeds in various concentrations of H₂O₂ resulted in enhanced stomatal density, and increased length and histological components of leaves. Those changes were positive since plants of H₂O₂-soaked seeds had higher dry weight [32] (Jafariyan and Zarea, 2016). Again, the higher doses (> 1%) of H₂O₂ probably caused toxic effects on the seedling dry matter and so, it became less, even than those of the unprimed and the hydro-primed seeds. Furthermore, Kaya et al. (2006) [28] experienced that priming increased seedling dry matter of sunflower under drought stress condition.

Table 2. Effect of priming seed with H_2O_2 on shoot, root and seedling dry matter of okra

| Treatments | Shoot dry matter | | | Root dry matter | | | Seedling dry matter | | |
|-----------------------|------------------|--------|---------|-----------------|---------|--------|---------------------|---------|---------|
| | 10 DAD | 20 DAD | 30 DAD | 10 DAD | 20 DAD | 30 DAD | 10 DAD | 20 DAD | 30 DAD |
| T ₁ | 6.01ab | 8.22ab | 18.20ab | 0.79bc | 1.46ab | 4.13ab | 6.80bc | 9.68ab | 22.11b |
| T ₂ | 6.88a | 8.63ab | 18.50ab | 0.76bc | 1.30bc | 3.79ab | 7.65ab | 9.933ab | 22.35b |
| T ₃ | 6.50ab | 8.68ab | 18.55ab | 0.86b | 1.37a-c | 3.97ab | 7.37abc | 10.15ab | 22.52b |
| T ₄ | 6.93a | 9.16a | 19.64a | 1.14a | 1.64a | 4.46a | 8.08a | 10.82a | 24.11a |
| T ₅ | 6.450ab | 7.66bc | 17.55bc | 0.84bc | 1.29bc | 3.83ab | 7.29a-c | 9.29b | 21.37bc |
| T ₆ | 6.57ab | 7.89b | 16.28cd | 0.86b | 1.31bc | 3.94ab | 7.44a-c | 9.55b | 20.2cd |
| T ₇ | 6.53ab | 7.80bc | 16.64cd | 0.78bc | 1.34bc | 3.69b | 7.31a-c | 9.15b | 20.33cd |
| T ₈ | 5.74b | 6.75c | 15.99d | 0.64c | 1.15c | 3.63b | 6.36c | 7.90c | 19.62d |
| LSD _(0.05) | 1.015 | 1.042 | 1.368 | 0.191 | 0.271 | 0.626 | 1.001 | 1.140 | 1.436 |
| CV % | 8.97 | 7.35 | 4.42 | 13.23 | 11.28 | 9.09 | 7.85 | 6.81 | 3.80 |



The figures with different letters differ among themselves at the 5% level of probability.

T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%)

3.5 Number of secondary roots and true leaves

The number of secondary roots and true leaves varied appreciably (P≤0.05) due to the effects of H₂O₂ concentrations at all the three DADs (Table 3). At the 10, 20 and 30 DAD, the highest number of secondary roots was recorded in T₄ (8.59, 14.78 and 20.30) but the lowest number was counted in T₈ (7.02, 11.66 and 17.86). At the 10, 20 and 30 DAD, the maximum number of true leaves was recorded from T₄ (0.91, 1.80 and 3.23) while the minimum was found in T₈ (0.80, 1.36 and 2.93). Again, the higher doses (over 1%) of H₂O₂ probably caused harmful

effects for the number of secondary roots and true leaves and so, those were fewer in number, even than the unprimed and the hydro-primed seeds. However, when the concentrations of H₂O₂ came up to 5mM, it played an opposite role to inhibit the growth of adventitious roots and seriously damaged those [33] (Deng et al., 2012). Similar types of result were also reported by [26] Lima (2017) in bitter gourd and bottle gourd where the highest number of secondary roots and true leaves was at 1.5% concentration of H₂O₂. Again, [25] Mustafa (2017) recorded the number of secondary roots (>1cm) and true leaves per seedling of cucumber, swampcabbage, radish and Indian spinach seeds primed with 1% H₂O₂. But the excessive accumulation of H₂O₂ leads to cellular oxidative damage and even programmed death [34-35] (Levine et al., 1994 and Prasad et al., 1994) and thus becomes poisonous for seedlings.

Table 3. Effect of priming seedwith H₂O₂ on number of secondary root and true leaves of okra

| Treatments | Number of secondary roots | | | Number of true leaves | | |
|-----------------------|---------------------------|---------|---------|-----------------------|---------|---------|
| | 10 days | 20 days | 30 days | 10 days | 20 days | 30 days |
| T ₁ | 7.40ab | 12.83bc | 19.09ab | 0.86a-c | 1.56bc | 3.057a |
| T ₂ | 6.56b | 14.57a | 18.08b | 0.90ab | 1.62ab | 3.167a |
| T ₃ | 7.51ab | 12.57bc | 18.86ab | 0.88a-c | 1.60ab | 3.143a |
| T ₄ | 8.59a | 14.78a | 20.30a | 0.91a | 1.80a | 3.233a |
| T ₅ | 7.15ab | 13.47ab | 18.07b | 0.83a-c | 1.63ab | 3.077a |
| T ₆ | 7.63ab | 12.06bc | 18.52b | 0.81bc | 1.65ab | 3.087a |
| T ₇ | 8.43a | 13.16b | 18.61b | 0.88a-c | 1.63ab | 2.947a |
| T ₈ | 7.02ab | 11.66c | 17.86b | 0.80c | 1.36c | 2.930a |
| LSD _(0.05) | 1.478 | 1.310 | 1.410 | 0.078 | 0.207 | 0.282 |
| CV % | 11.19 | 5.70 | 4.31 | 5.12 | 7.37 | 5.23 |

The figures with different letters differ among themselves at the 5% level of probability.

T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%)

3.6 Seedling vigor index

The seedling vigor index differed significantly (P≤0.05) due to the different concentrations of H₂O₂ at all the three DADs

(Table 4). At the 10, 20 and 30 DAD, the highest seedling vigor index was recorded from T₄ (958.3, 1882 and 2270, respectively) while the least value was recorded from T₈ (706.8, 1882 and 1824, respectively). [25] Mustafa (2017) also recorded the maximum seedling vigor index of swamp cabbage, radish and Indian spinach seeds primed with 1%, H₂O₂. [26] Lima (2017) and [24] Nandi et al. (2017) found similar findings on seedling vigor index.

Table 4. Effect of seed priming with H₂O₂ on Seedling vigor index,relative growth rate and root shoot ratio of okra

| Treatments | Seedling vigor index | | | Relative growth rate | | Root shoot ratio | | |
|-----------------------|----------------------|--------|---------|----------------------|--------|------------------|--------|--------|
| | 10 DAD | 20 DAD | 30 DAD | 20 DAD | 30 DAD | 10 DAD | 20 DAD | 30 DAD |
| T ₁ | 787.8b-d | 1527cd | 1938b-d | 0.033a | 0.076a | 0.130a | 0.173a | 0.220a |
| T ₂ | 758.8cd | 1556cd | 1886c-e | 0.020a | 0.076a | 0.110a | 0.143a | 0.200a |
| T ₃ | 814.3b | 1585c | 2028b | 0.016a | 0.076a | 0.133a | 0.153a | 0.210a |
| T ₄ | 958.3a | 1882a | 2270a | 0.026a | 0.076a | 0.163a | 0.180a | 0.223a |
| T ₅ | 807.8bc | 1785b | 1977bc | 0.016a | 0.076a | 0.126a | 0.186a | 0.213a |
| T ₆ | 812.0b | 1503d | 1974bc | 0.020a | 0.070a | 0.126a | 0.176a | 0.236a |
| T ₇ | 742.5de | 1523cd | 1869de | 0.020a | 0.073a | 0.120a | 0.166a | 0.216a |
| T ₈ | 706.8e | 1413e | 1824e | 0.016a | 0.070a | 0.106a | 0.176a | 0.220a |
| LSD _(0.05) | 49.02 | 69.88 | 87.85 | 0.0553 | 0.0553 | 0.0553 | 0.055 | 0.055 |



| | | | | | | | | |
|------|------|------|------|-------|-------|-------|-------|-------|
| CV % | 3.51 | 2.50 | 2.55 | 40.43 | 14.74 | 17.21 | 13.15 | 10.98 |
|------|------|------|------|-------|-------|-------|-------|-------|

The figures with different letters differ among themselves at the 5% level of probability.

3.7 Relative growth rate (RGR)

There was no significant variation in the relative growth rate among the treatments at 20 and 30 DADs (Table 4). numerically, the maximum relative growth rate was in T₁ (0.033 and 0.076), the minimum relative growth rate was found in T₈ (0.016 and 0.070) at the 20 and 30 DAD. While working with, [25] Mustafa (2017) found the highest relative growth rate at 3% (0.22) in cucumber and Indian spinach seeds. [26] Lima (2017) found relative growth rate in control/no priming (0.11) in bitter gourd seed and bottle gourd seeds.

3.8 Root shoot ratio

There was statistically non-significant variation in root shoot ratio (dry weight basis) among the different treatments at 10, 20 and 30 DAD (Table 4). [26] Lima (2017) and [25] Mustafa (2017) found the maximum root shoot ratio treated with 1% H₂O₂.

IV. CONCLUSIONS

So, from the results of the experiment, it can be concluded that H₂O₂ had optimistic effects on the seedling qualities. Again, among the six concentrations of the H₂O₂, 1% was the most effective one for okra seeds. Above those concentrations, others were somewhat toxic as those hindered a lot of the parameters evaluated.

V. REFERENCES

- [1]. Kochhar SL (2004). Economic Botany in the tropics. Macmillan India Limited, Daryaganj, New Delhi, 604.
- [2]. Sharma DK, Jain VK, Rajni J and Nandini S. (2013). Post-harvest study of okra (*Abelmoschus esculentus* (L.) Moench) fruits and phytopathological effect of associated microflora. *International Journal of Innovative Research and Review*, 1 (1), 27-34.
- [3]. Sharma DK, Jain VK, Jain R and Sharma N. (2013). Effects of microflora associated with okra (*Abelmoschus esculentus* L.) Moench seeds and their phytopathological effects. *Cibtech Journal of Microbiology*, 2(2) 39-44.
- [4]. Sathish KD, Eswar TD, Praveen KA, Ashok KK, BramhaSrinivasa RD and Ramarao N. (2013). A review on: *Abelmoschus esculentus* (Okra). *International Research Journal of Pharma Applied Sciences*, 3(4), 129-132.
- [5]. Priya S, Varun C, Tiwari BK, Chauhan SS, Sobita S, Bilal S and Abidi AB. (2014). An overview on okra (*Abelmoschus esculentus*) and its importance as a T₁ (control/no priming), T₂ (hydro-priming in plain H₂O) and other six aquatic solutions of H₂O₂, T₃ (0.5%), T₄ (1.0%), T₅ (1.5%), T₆ (2.0%), T₇ (2.5%) and T₈ (3.0%) nutritive vegetable in the world. *IJPBS*, 4(2), 227-233.
- [6]. Lokesh. (2017). Nutritional and pharmaceutical potentials of okra (*Abelmoschus esculentus*) plant and its biotic stresses -An Overview. *International Journal of Pure Applied Biosciences*, 5(4), 1890-1907.
- [7]. Kranner, I., Chen, H. Y., Pritchard, H. W., Pearce, S. R. and Birtic, S. 2011. Seed ageing correlates with inter-nucleosomal DNA fragmentation and loss of RNA integrity. *Plant Growth Regul.* 63: 63–72.
- [8]. Schopfer, P., Plachy, C. and Frahry, G. 2001. Release of reactive oxygen intermediates (Superoxide radicals, H₂O₂ and hydroxyl radicals) and peroxides in germinating radish seeds controlled by light, gibberellins and abscisic acid. *Pl. Physiol.* 125: 1591-1602.
- [9]. Weitbrecht, K., Müller, K. and Leubner-Metzger, G. 2011. First of the mark: early seed germination. *J. Exp. Bot.* 62: 3289–3309.
- [10]. Taylor, A. G. and Harman, G. E. 1998. Concepts and technologies of selected seed treatments. *Annu Rev Phytopathol.* 28: 321–39.
- [11]. Bradford, K. J. 1986. Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *Hort. Sci.* 21: 1105-1112.
- [12]. Finnerty, T. L., Zajicek, J. M. and Hussey, M. A. 1992. Use of seed priming to bypass stratification requirements of three *Aquilegia* species. *Hort Science.* 27: 310-313.
- [13]. Coolbear, P. and McGill, C. R. 1990. Effect of low temperature pre-sowing treatment on the germination of tomato seeds under temperature and osmotic stress. *Scientia Horticulturae* 44: 43-54.
- [14]. Knypl, J. S. and Khan, A. A. 1981. Osmo-conditioning of soybean seeds to improve performance at suboptimal temperatures. *Agronomy Journal.* 73: 112-116.
- [15]. Hardegree, S. P. and Emmerich, W. E. 1992. Seed germination response of four south-western range grasses to equilibration at sub-germination matrix potentials. *Agronomy Journal.* 84: 994-998.
- [16]. Toselli, M. E. and Casenave, E. C. 2003. Water content and the effectiveness of hydro and osmotic priming of cotton seeds. *Seed Sci. and Tech.* 31: 727-735.
- [17]. Black, M., Bewley, J. D., Halmer, P. 2006. *The Encyclopedia of Seeds: Science, Technology and Uses.* Wallingford: CABI. 5-6



- [18]. Jann, R. C. and Amen, R. D. 1997. What is germination? In: Khan A. A. (Ed.). The physiology and biochemistry of seed dormancy and germination, North Holland Publishing, Amsterdam, 7-28.
- [19]. Jia, W., Wang, Y., Zhang, S. and Zhang, J. 2002. Salt-stress-induced ABA accumulation is more sensitively triggered in roots than in shoots. *J. Expt. Bot.* 53: 2201-2206.
- [20]. Ogawa, K. and Masaki, I. 2001. A Mechanism for promoting the germination of *Zinnia elegans* seeds by hydrogen peroxide. *Plant cell physiol.* 42(3): 286-291.
- [21]. Patade, V. Y., Maya, K. and Zakwan, A. 2012. Seed priming mediated germination improvement and tolerance to subsequent exposure to cold and salt stresses in capsicum. *Res. J. Seed Sci.* 4: 125-136.
- [22]. Copeland, L.O. and McDonald, M. B. 1985. Principles of seed science and technology. Macmillan Publishing Co. USA. P.3-5.
- [23]. John, R. and Duval. 2000. Treatment with hydrogen peroxide and seed coat removal or clipping improves germination of 'Genesis' triploid watermelon. *Hort. Sc.* 35(1): 85-86.
- [24]. Nandi, M., Pervez, Z., Alam, S., Islam, S. and Mahmud, R. 2017. Effect of hydrogen peroxide treatment on health and quality of chilli seed. *Int. J. Pl. Pathol.* 8: 8-13.
- [25]. Mustafa, H. 2017. Improving the seedling qualities of some vegetables priming with H_2O_2 . An M. S. thesis, Dept. of Horticulture, Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur. pp. 1-89.
- [26]. Lima, I. J. 2017. Enhancing the seedling qualities of some vegetables priming with H_2O_2 . An M. S. thesis, Dept. of Horticulture, Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur. pp. 1-67.
- [27]. Iqbal, T.M.T., Bahadur, M.M., Kabir, M.A., Hasan, M.A. and Majumder, D.N.A. 2001. Improvement of okra seed quality by pre-soaking in H_2O_2 solution. *Pakistan J. of Bio. Sci.* 4(6): 639-641.
- [28]. Kaya, D., M., Okçu, G., Atak, M., Çikili, Y. and Kolsarici, Ö. 2006. Seed treatment to overcome salt and drought stresses during germination in sunflower. *Europ. J. of Agrn.* 24: 291-295.
- [29]. Barba-Espin, G., Diaz-Vivancos, P., Clemente-Moreno, M.J., Albacete, A., Faize, L., Faize, M., F. Pérez-Alfoceaand, F. and Hernández, J.A. 2010. Interaction between hydrogen peroxide and plant hormones during germination and the early growth of pea seedlings. *Pl., Cell and Environ.* 33: 981-994.
- [30]. Ogiwara, H. and Terashima, K. 2001. A varietal difference in coleoptile growth is correlated with seedling establishment of direct seeded rice in submerged field under low temperature conditions. *Plant Prod. Sci.* 4: 166-172.
- [31]. León-Vargas, Julio, Guevara-Acevedo, Luis, Cervantes-ortíz, Francisco, Rodríguezmer-cado, Daniel, Cisneros-López, Hugo and Adrío-Enríquez and Enrique. 2016. Effect of hydrogen peroxide on broccoli seedlings. *Ecorfan J. Ecuador*, 3-5:7-13.
- [32]. Rahman, M.M., Rahman, M.M. and Islam, M.N. 1997. Effect of H_2O_2 pre-soaking on germination and early growth of groundnut seeds. *Bangla. J. Seed Science and Tech.* 1:63-68.
- [33]. Jafariyan, T. and Zarea, M. J. 2016. Hydrogen peroxide affects plant growth promoting effects of *Azospirillum*. *J. Crop Sci. Biotech.* 19(2): 167-175.
- [34]. Deng, X. P., Cheng, Y. J., Wu, X. B., Kwak, S. S., Chen, W. and Eneji, A. E. 2012. Exogenous hydrogen peroxide positively influences root growth and metabolism in leaves of sweet potato seedlings. *6(11): 1572-1578.*
- [35]. Levine, A., Tenhaken, R., Dixon R. and Lamb, C. 1994. H_2O_2 from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell.* 7: 583-593.
- [36]. Prasad, T. K., Anderson, M. D., Martin, B. A. and Stewart, C. R. 1994. Evidence for chilling-induced oxidative stress in maize seedlings and a regulatory role for hydrogen-peroxide. *Plant Cell.* 6: 65-74.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143